

The Multi-Layer Multi-Configuration Time-Dependent Hartree Method for Bosons: Theory and Applications

Lushuai Cao^{a,1,2,*} Sven Krönke^{a,1,2,†} Oriol Vendrell^{3,2,‡} and Peter Schmelcher^{1,2,§}

¹*Zentrum für Optische Quantentechnologien, Universität Hamburg,
Luruper Chaussee 149, D-22761 Hamburg, Germany*

²*The Hamburg Centre for Ultrafast Imaging,
Luruper Chaussee 149, D-22761 Hamburg, Germany*

³*Center for Free-Electron Laser Science, DESY,
Notkestrasse 85, D-22607 Hamburg, Germany*

Abstract

In this work, we develop and apply the multi-layer multi-configuration time-dependent Hartree method for bosons (ML-MCTDHB), which represents a unique tool for investigating the non-equilibrium dynamics of multi-species bosonic systems in arbitrary dimensions. Being an ab initio method for solving the time-dependent Schrödinger equation, ML-MCTDHB takes all correlations into account. The multi-layer feature of ML-MCTDHB allows for tailoring the wave function ansatz in order to describe intra- and inter-species correlations accurately and efficiently. To show the beneficial scaling and the efficiency of the method, we explore the correlated dynamics of three species tunneling in a double well trap. Here we demonstrate and analyze in detail the build up of inter- and intra-species correlations in the course of the quantum dynamics as well as signatures of equilibration. Furthermore, we show that the presence of the third species which ultra-weakly attracts bosons of the two other species can reduce the correlations between those two species.

PACS numbers: 03.75.Kk, 05.30.Jp, 03.65.-w, 31.15.-p

^a These two authors contributed equally.

^{*} lcao@physnet.uni-hamburg.de

[†] Sven.Kroenke@physnet.uni-hamburg.de

[‡] oriol.vendrell@cfel.de

[§] pschmelc@physnet.uni-hamburg.de

I. INTRODUCTION

Due to the high degree of controllability and isolatedness, ensembles of ultracold atoms have been attracting much attention for almost two decades. Whereas the first experiments with ultracold bosons were situated in parameter regimes which allow for a mean field description by means of the famous Gross Pitaevskii equation [1–3], nowadays experimental techniques enable us to also study the effects of strong interactions leading to highly correlated states [4]. Furthermore, different trapping geometries resulting in lower dimensional ensembles as well as effects due to the presence of other components (e.g. [5]) are in the focus. In particular, it has become feasible to trap and manipulate ensembles of only a few atoms (e.g. [6, 7]), which allows to explore the impact of the particle number on various effects. The quest for a description of the non-equilibrium quantum dynamics beyond the mean field approach, i.e. by taking many-body correlations into account, demands new theoretical tools.

Simulating the quantum dynamics of an interacting N particle system, however, is a tough task in general due to the exponential scaling of the state space: If, for instance, one allows each particle to occupy M single particle states, the many-body Hilbert space turns out to be M^N dimensional for a system of distinguishable particles. When going to higher particle numbers, one, hence, has to truncate the Hilbert space further. Moreover, this scaling problem has a serious consequence for simulating *dynamics*: In the course of time, the system state will move through different subspaces of the total Hilbert space in general. If one now tries to expand the system state with respect to a time-independent basis, many basis states will be needed in order to span the relevant subspace for the whole propagation time. So for reducing the number of basis vectors, one should better employ a time-dependent, moving basis, in which the instantaneous system state can be optimally represented.

In the multi-configuration time-dependent Hartree method (MCTDH), such a co-moving basis is provided by time-dependent Hartree products [8, 9]. The equations of motion for the single particle functions (SPFs) building up the time-dependent Hartree products as well as for the expansion coefficients are obtained by means of the Dirac-Frenkel variation principle, which ensures a variationally optimal representation of the total wave function at any instant. Using Hartree products as the many-body basis states, MCTDH is well suited

for dealing with distinguishable particles. Originally developed in the context of quantum molecular dynamics, MCTDH has successfully been applied to an enormous diversity of problems concerning the vibrational and wave packet dynamics of molecules (cf. [10, 11] and reference therein). Moreover, MCTDH turned out to be also powerful for simulating strongly correlated few-body systems of ultracold, indistinguishable bosonic atoms in traps, see e.g. [12, 13].

MCTDH has been extended for dealing with higher particle numbers in several ways: In multi-layer MCTDH (ML-MCTDH), degrees of freedom are combined to higher-dimensional ones. The MCTDH expansion scheme is then applied to these combined modes and their constituting degrees of freedom in a cascade. The ML-MCTDH theory has been presented and implemented for the first time in [14]. Afterwards, a recursive formulation of the ML-MCTDH theory was derived, which allows for a fast and convenient treatment of arbitrarily complex ML-MCTDH expansion schemes [15]. The ML-MCTDH method is particularly fruitful for scenarios where there are strong correlations within certain subsystems while the inter-subsystem correlations are relatively weak [16–22]. Especially, many system-bath problems are of this type. In such a situation, the necessary number of time-dependent basis functions can be reduced by combining the strongly correlated degrees of freedom in an ML-MCTDH expansion. As MCTDH, ML-MCTDH is based on expansions in terms of Hartree products and, thus, ignores possible particle exchange symmetries of the system.

In contrast to this, the MCDTH method for fermions (MCTDHF) and for bosons (MCTDHB) reduce the number of expansion coefficients by expanding the total wave function in terms of Slater determinants and permanents [23–26], respectively. MCTDHF has been applied in the context of quantum chemistry as well as for molecules in strong laser fields [27–30]. The crossover from the Gross-Pitaevskii to the many-body regime has been studied for several ultracold bosonic systems by means of MCTDHB (e.g. [31, 32]). There is even a formalism, which unifies MCTDHB and MCTDHF for dealing with Bose-Bose, Bose-Fermi and Fermi-Fermi mixtures [33]. Furthermore, an extension for considering particle conversion in a multi-species system has been developed [34].

ML-MCTDH features a high flexibility with respect to the wave function expansion: Any way of combining degrees of freedom to higher dimensional ones in a cascade is allowed and leads to a specific wave function ansatz. As, in the case of indistinguishable particles, many of these mode-combination schemes do not reflect the indistinguishability of the particles,

the particle exchange (anti-) symmetry can, in general, not be incorporated in ML-MCTDH, making ML-MCTDH incompatible with MCTDHB and MCTDHF. There is, however, an exception: In [35], a second quantization representation of ML-MCDTH (ML-MCTDH-SQR) has been derived, which has been applied to charge transport problems [36, 37], for example. The ML-MCTDH-SQR method is based on the factorization of the Fock space for m single particle modes into m Fock spaces corresponding to just one single particle mode each, which leads to the product structure required by ML-MCTDH. In this way, ML-MCTDH-SQR benefits from the flexible cascade scheme of ML-MCTDH while keeping the SPF, constituting the Fock states, time-independent.

For investigating complex bosonic systems, a new ab initio method, called multi-layer multi-configuration time-dependent Hartree method for bosons (ML-MCTDHB), is derived and applied to explore novel quantum dynamics in this work. This method provides a new symbiosis of MCTDHB and ML-MCTDH, which in particular differs from ML-MCTDH-SQR in the way that it is based on time-dependent SPFs. To face the challenge of simulating the non-equilibrium dynamics of complex bosonic systems without discarding any correlations, we derive two basic classes of ML-MCTDHB wave function ansatzes: (i) a multi-layer expansion with respect to the different bosonic species for optimally taking account inter- and intra-species correlations and (ii) a particle multi-layer expansion in which the single particle functions are further expanded in the MCTDH form. The latter allows for an optimal wave function ansatz when dealing with trap geometries in higher dimensions or internal degrees of freedom. In both classes of expansion schemes, the bosonic exchange symmetry is exactly taken into account, which improves the accuracy as well as the efficiency of the method. Furthermore, any combination of the expansion schemes (i) and (ii) is possible resulting in a highly flexible ab initio approach to systems with - in principle - arbitrary many species in arbitrary dimensions.

This paper is organized as follows: Firstly, we will review the methods ML-MCTDH and MCTDHB in sections II A and II B. The wave function expansion of ML-MCTDHB is consecutively introduced in section II C 1 and afterwards the derivation principle for the equations of motion as well as the results for multi- and single-species systems are presented in sections II C 2, II C 3 and II C 4, respectively. In section III, ML-MCTDHB is applied to several tunneling scenarios, which show interesting correlation effects and at the same time illustrate how the ML-MCTDHB wave function ansatz can be adapted to capture

the relevant physics. We conclude with a summary, discussing also an extension of ML-MCTDHB to fermionic species in section IV. Finally, we would like to note that the following reviews of the existing methods ML-MCTDH and MCTDHB are indispensable in order to be self-contained, to embed ML-MCTDHB into the existing methods, to give a consistent definition of the layer structure in connection with particle statistics and in order to be able to present ML-MCTDHB in its full generality.

II. THEORY

A. ML-MCTDH

Firstly, let us summarize the MCTDH and ML-MCTDH approaches [8–11, 14–16, 38]. We will review these two methods in such a way that an extension to the subject of this work, ML-MCTDHB, can be achieved smoothly. At the first place, we will introduce the wave function expansion schemes for both MCTDH and ML-MCTDH. Afterwards, the equations of motion are discussed.

Let us consider the general situation of a system consisting of N physical degrees of freedom (DOF) $(x_1, x_2, \dots, x_N)$, where the DOF x_κ can correspond to a position, momentum or internal degree of freedom. Furthermore, N does not need to correspond to the number of physical particles but can take internal degrees of freedom as well as the dimensionality of the system into account. For instance, a system of N physical particles in three-dimensional space can either be modeled by $3N$ DOF, $(x_1, x_2, \dots, x_{3N})$, or by N three-dimensional DOF, $(x_1, x_2, \dots, x_N)$. To each physical DOF κ , we then associate a set of orthonormal time-dependent single particle functions (SPFs) $\{\phi_n^\kappa(x_\kappa), n = 1, \dots, m_\kappa\}$. In MCTDH, the state vector of the system is expanded with respect to the (truncated) basis of Hartree products constituted by the SPFs of each DOF:

$$|\Psi(t)\rangle = \sum_{i_1, i_2, \dots, i_N} A_{i_1, i_2, \dots, i_N}(t) |\phi_{i_1}^1(t)\rangle |\phi_{i_2}^2(t)\rangle \dots |\phi_{i_N}^N(t)\rangle \equiv \sum_I A_I |I\rangle. \quad (1)$$

To step from MCTDH to ML-MCTDH, we first perform a mode combination of the N physical DOF and group them into N_{P-1} so-called logical DOF such that the first n_1 physical DOF are combined to the logical DOF x_1^{P-1} , the next n_2 physical DOF are combined to the

logical DOF x_2^{P-1} , and so on:

$$\begin{aligned}
x_1^{P-1} &= x_1^{P-1}(x_1, x_2, \dots, x_{n_1}), \\
x_2^{P-1} &= x_2^{P-1}(x_{n_1+1}, x_{n_1+2}, \dots, x_{n_1+n_2}), \\
&\vdots \\
x_{N_{P-1}}^{P-1} &= x_{N_{P-1}}^{P-1}(x_{N-n_{N_P}+1}, x_{N-n_{N_P}+2}, \dots, x_N).
\end{aligned} \tag{2}$$

The physical DOF $\{x_i\}$ are said to form the physical layer in ML-MCTDH, and the logical DOF $\{x_i^{P-1}\}$ constitute the first upper layer above the physical layer, where the superscript $P-1$ denotes that these DOF are one layer above the physical layer. We can continue and combine now the N_{P-1} logical DOF $\{x_i^{P-1}\}$ to generate a new set of N_{P-2} ($N_{P-2} < N_{P-1}$) logical DOF $\{x_i^{P-2} | i \in [1, N_{P-2}]\}$, which form the second upper layer above the physical layer. By continuing this mode-combination scheme until reaching a layer containing only one logical DOF, we complete the construction of the so-called tree structure in ML-MCTDH. Generally speaking, the tree structure of ML-MCTDH contains different layers of DOF, and the DOF on each layer α are generated via mode combination of the corresponding DOF on the lower lying layer $\alpha+1$.

Similarly to the physical DOF, we also associate a set of orthonormal time-dependent SPFs $\{|\phi_n^{\alpha;\kappa}\rangle\}$ to each logical DOF x_κ^α . Please note that, despite of the term ‘‘SPF’’, the state $|\phi_n^{\alpha;\kappa}\rangle$ can correspond e.g. only to the x - and y -coordinate of a particle in three-dimensional coordinate space or to the x -coordinates of four different particles, depending on the mode-combination scheme.

Each of these SPFs is expanded with respect to Hartree products made of the SPFs of those DOF $x_{\kappa'}^{\alpha+1}$ which constitute the logical DOF x_κ^α :

$$|\phi_n^{\alpha;\kappa}\rangle = \sum_{i_1, i_2, \dots, i_m} A_{n; i_1, i_2, \dots, i_m}^{\alpha;\kappa} |\phi_{i_1}^{\alpha+1;\kappa_1}\rangle |\phi_{i_2}^{\alpha+1;\kappa_2}\rangle \dots |\phi_{i_m}^{\alpha+1;\kappa_m}\rangle \equiv \sum_I A_{n; I}^{\alpha;\kappa} |I^{\alpha;\kappa}\rangle. \tag{3}$$

We note that the time-dependencies are omitted in the notation. Here and from now on, the ingredients of all formulas have to be regarded as being time-dependent, unless stated otherwise.

Especially, the single DOF of the top layer possesses only a single SPF, which turns out to be the state vector $|\Psi\rangle$ of the system and is expanded in terms of Hartree products made

of the SPFs of the second layer:

$$|\Psi\rangle = \sum_I A_I^1 |I^1\rangle. \quad (4)$$

Equations (3) and (4) indicate that each SPF is “locally” expanded in the MCTDH manner (1), in terms of Hartree products made of the SPFs of the lower layer. Finally, the ML-MCTDH wave function representation is completed by expanding the SPFs of the physical DOF with respect to some given time-independent basis function $|x_\kappa\rangle$, which one usually calls *primitive basis functions* (PBFs):

$$|\phi_n^\kappa\rangle \equiv |\phi_n^{P;\kappa}\rangle = \sum_{x_\kappa} \phi_n^\kappa(x_\kappa) |x_\kappa\rangle = \sum_{x_\kappa} A_{n;x_\kappa}^{P;\kappa} |x_\kappa\rangle. \quad (5)$$

The physical DOF x_κ can be high-dimensional, for instance, x_κ might contain the x -, y -, and z -position degrees of freedom for the case of particles in three dimensional coordinate space. In this example, the SPFs of the DOF x_κ would be expanded in terms of three-dimensional PBFs $|x_\kappa\rangle \equiv |\vec{r}_\kappa\rangle$, which is known as mode combination in MCTDH (cf. [8]). We will call such a mode combination *primitive mode combination* in order to distinguish it from the mode combinations in the other layers of a ML-MCTDH wave function expansion. Finally we remark that having the sets of PBFs fixed, the evolution of the system turns out to be completely represented by the time evolution of the expansion coefficients of each DOF, $A_{i;I}^{\alpha;\kappa}$.

Before we state the equations of motions for those coefficients $A_{i;I}^{\alpha;\kappa}$, we briefly review the diagrammatic representation of the wave function expansion (3,4,5) introduced by Manthe [15]: Performing the MCTDH expansion layer by layer in a cascade naturally leads to a visualization in terms of a tree diagram. In such a tree diagram, every node is identified with a logical or physical DOF x_κ^α . Its number of teeth represents how many SPFs $|\phi_i^{\alpha;\kappa}\rangle$ are provided for this DOF. Then each link between two nodes indicates the summation over an index in the expansion (3,4,5). Finally, a rectangular box stands for a set of PBFs and the number of its teeth coincides with the number of given PBFs. Figure 1 illustrates this diagrammatic notation by an example of two particles in two-dimensional coordinate space: The physical layer consists of four DOF corresponding to the two spatial directions for each particle. In the second layer, two DOF each are combined to one logical DOF each, representing the two physical particles. In the following we will equivalently refer to the DOF as nodes. This will enable us to address the DOF, which turn out to be relevant for

a node under consideration, by the pictorial terms *parent* and *child node*, respectively. At this step, it is clear that MCTDH could be seen as a special case of ML-MCTDH, which contains only two layers: the top layer of $|\Psi(t)\rangle$ and the primitive layer of physical DOF. In this way, we will mainly focus on the equations of motion of ML-MCTDH, and those of MCTDH are naturally included.




Figure 1: Diagrammatic representation of a ML-MCTDH wave function expansion for a system of two physical particles, which correspond to the nodes of the second layer. Indicated by their teeth, four and two SPFs are provided for the first and second physical particle, respectively. According to the third layer, each of these SPFs is unraveled by Hartree products of two one-dimensional SPFs, i.e. orbitals of the x- and y-direction. Each of these one-dimensional orbitals is in turn represented by n_x and n_y , respectively, PBFs or, pictorially speaking, spatial gridpoints. For illustrating the notation, the expansion coefficients are written into the corresponding nodes.

We assume the Hamiltonian of the system to be given as a sum over products of single particle operators acting on each physical DOF, which is particularly suitable for ML-MCTDH as it allows for a recursive formulation of the theory layer by layer:

$$\hat{H} = \sum_{\nu} C_{\nu} \prod_{\kappa=1}^N \hat{h}_{\nu}^{\kappa} \equiv \sum_{\nu} C_{\nu} \hat{H}_{\nu}. \quad (6)$$

Here, $\hat{h}_{\nu}^{\kappa}$ denotes a single particle operator¹ acting on the physical DOF x_{κ} . Some interaction

¹ Please note that the term “single particle operator” does not need to refer to a physical particle but refers

potentials can be brought analytically into the desired product form (6), see the harmonic interaction $v(x_1 - x_2) \propto (x_1 - x_2)^2$, for example. Moreover, such a decomposition of a many-body operator into products of single particle operators can always be realized by the so-called POTFIT algorithm, which essentially solves N eigenvalue problems [39, 40]. For a Hamiltonian with up to F -body operators, the number of ν terms in (6) in general equals the product of the dimensions of the Hilbert spaces corresponding to the respective F physical DOF but one of these. The ν term summation, however, can be truncated in a controlled way, and often, only a few ν terms are necessary for a fair representation of the total Hamiltonian. In this case, the product form (6) can improve the efficiency because high-dimensional integrals decompose into low-dimensional ones when calculating matrix elements. We remark that the Hamiltonian $\hat{H}$ may be time-dependent in ML-MCDTH.

Since every layer in itself provides a complete description of the system, the Hamiltonian can be also described equivalently by the SPFs of each layer: For the α layer, the single particle operators $\hat{h}_\nu^\kappa$ can be grouped according to the mode-combination scheme up to the layer α , which leads to the α -layer representation of the Hamiltonian $\hat{H}$:

$$\hat{H} = \sum_{\nu} C_{\nu} \prod_{\kappa=1}^{N_{\alpha}} \hat{h}_{\nu}^{\alpha;\kappa} \equiv \sum_{\nu} C_{\nu} \hat{H}_{\nu}^{\alpha}. \quad (7)$$

Here, $\hat{h}_{\nu}^{\alpha;\kappa}$ denotes an operator exclusively acting on the logical DOF x_{κ}^{α} . Hence, it can be expressed in terms of the SPFs of this DOF as well as the operators of the layer $\alpha + 1$, whose product equals $\hat{h}_{\nu}^{\alpha;\kappa}$:

$$[h_{\nu}^{\alpha;\kappa}]_{nm} \equiv \langle \phi_n^{\alpha;\kappa} | \hat{h}_{\nu}^{\alpha;\kappa} | \phi_m^{\alpha;\kappa} \rangle = \sum_{I,J} (A_{n;I}^{\alpha;\kappa})^* A_{m;J}^{\alpha;\kappa} \langle I^{\alpha;\kappa} | \prod_{i=\kappa_1}^{\kappa_n} \hat{h}_{\nu}^{\alpha+1;i} | J^{\alpha;\kappa} \rangle, \quad (8)$$

where the DOF $(\alpha; \kappa)$ is a combination of the DOF $(\alpha + 1, \kappa_1)$ to $(\alpha + 1, \kappa_n)$.

A useful concept for the equations of motion is the single hole function (SHF) of the node $(\alpha; \kappa)$, $|\Psi_i^{\alpha;\kappa}\rangle$, which is given by the projection of the total wave function onto the i -th SPF of the node $(\alpha; \kappa)$:

$$|\Psi_i^{\alpha;\kappa}\rangle \equiv \langle \phi_i^{\alpha;\kappa} | \Psi \rangle. \quad (9)$$

The SHF of a node $(\alpha; \kappa)$ can be calculated by expanding the total wave function in terms of the SPFs of the layer α and performing the projection (9). A recursive formula for obtaining the SHFs layer by layer is stated in [15].

to a physical DOF.

Now we can substitute the ansatz of $|\Psi\rangle$ to the Dirac-Frenkel variational principle [41, 42]:

$$\langle \delta\Psi | (i\partial_t - \hat{H}) | \Psi \rangle = 0, \quad (10)$$

to obtain the equations of motion for the coefficients of each layer. The Dirac-Frenkel variational principle turns out to be equivalent to the Lagrangian and to the McLachlan's one [43, 44]. The latter essentially says: Propagate your ansatz $|\Psi\rangle$ according to $i\partial_t|\Psi\rangle = |\Theta\rangle$ with $|\Theta\rangle$ being of the ansatz form and minimizing $\|i|\Theta\rangle - \hat{H}|\Psi\rangle\|^2$. This ensures that we obtain a variationally optimal wave function within our class of ansatzes. Let us turn back to (10), which is easier to handle: The variation with respect to the top node coefficients, i.e. $\frac{\delta}{\delta(A_I^1)^*} \langle \Psi | (i\partial_t - \hat{H}) | \Psi \rangle = 0$, effectively gives the projection of the Schrödinger equation on the Hartree product $|I^1\rangle$. This results in the equation of motion for the top layer coefficients:

$$i\partial_t A_I^1 = \sum_J \langle I^1 | \hat{H} | J^1 \rangle A_J^1. \quad (11)$$

The variation with respect to a non-top node coefficient results in effectively projecting the Schrödinger equation on the SHF $|\Psi_n^{\alpha;\kappa}\rangle$ times the Hartree product $|I^{\alpha;\kappa}\rangle$. After some algebra, one obtains the equation of motion for the coefficients $A_{n,I}^{\alpha;\kappa}$:

$$\begin{aligned} i\partial_t A_{n,I}^{\alpha;\kappa} = \sum_\nu C_\nu \{ \sum_J \langle I^{\alpha;\kappa} | (1 - \hat{P}^{\alpha;\kappa}) \hat{h}_\nu^{\alpha;\kappa} | J^{\alpha;\kappa} \rangle \} \times \\ \times \{ \sum_m [(\rho^{\alpha;\kappa})^{-1} * \langle H_\nu^{\alpha;\kappa} \rangle]_{n,m} \} A_{m,J}^{\alpha;\kappa}. \end{aligned} \quad (12)$$

Here, $[(\rho^{\alpha;\kappa})^{-1} * \langle H_\nu^{\alpha;\kappa} \rangle]$ denotes the matrix product of the inverse² of the density matrix of DOF x_κ^α and the so-called mean field matrix of the ν -th Hamiltonian term with respect to x_κ^α : The density matrix $\rho^{\alpha;\kappa}$ is defined as the overlap of different SHFs of the DOF x_κ^α :

$$\rho_{n,m}^{\alpha;\kappa} := \langle \Psi_n^{\alpha;\kappa} | \Psi_m^{\alpha;\kappa} \rangle. \quad (13)$$

It turns out that $\rho_{n,m}^{\alpha;\kappa}$ equals the transposed of the physical reduced density matrix of the DOF x_κ^α , which one obtains by tracing out all DOF but x_κ^α in the many-body state $|\Psi\rangle\langle\Psi|$. The mean field matrix $\langle H_\nu^{\alpha;\kappa} \rangle_{n,m}$ is defined as the scalar factor of the mean field operator $\langle \Psi_n^{\alpha;\kappa} | \hat{H}_\nu^{\alpha;\kappa} | \Psi_m^{\alpha;\kappa} \rangle$ when making use of the product structure of $\hat{H}_\nu^\alpha$ (cf. (7)):

$$\hat{h}_\nu^{\alpha;\kappa} \langle H_\nu^{\alpha;\kappa} \rangle_{n,m} := \langle \Psi_n^{\alpha;\kappa} | \hat{H}_\nu^{\alpha;\kappa} | \Psi_m^{\alpha;\kappa} \rangle, \quad (14)$$

² As a density matrix can be singular and will be singular especially for converged MCTDH type simulations, one has to regularize the density matrix in (12), which can be done without affecting the total wave function (cf. [8]).

which can be evaluated by means of (8) in a recursive manner [15]. The operator $\hat{P}^{\alpha;\kappa}$ equals the projection operator onto the space spanned by the SPFs of the node $(\alpha;\kappa)$.

We note the following interpretation of the terms in (12): The operator $\hat{h}_\nu^{\alpha;\kappa}$ is responsible for the transition between different Hartree products $|J^{\alpha;\kappa}\rangle$, whereas the density and the mean field matrices scatter the SPF $|\phi_m^{\alpha;\kappa}\rangle$ to $|\phi_n^{\alpha;\kappa}\rangle$. Furthermore, it has to be pointed out that the equations of motion (11,12) are valid under the additional set of constraints:

$$\langle \partial_t \phi_i^{\alpha;\kappa} | \phi_j^{\alpha;\kappa} \rangle = 0, \quad (15)$$

which obviously ensures the orthonormality of the SPFs. In the following sections, we implicitly impose the constraint (15) on all occurring SPFs since such an additional condition is needed for obtaining unique equations of motions (cf. [8]).

Finally, we would like to emphasize that the flexible cascade of MCTDH expansions (3,4,5) leads to an optimal resolution of the relevant subspace of the many-body Hilbert space within the truncated basis given by the SPFs. Nevertheless, both MCTDH and ML-MCTDH are methods tailored for treating distinguishable DOF. In the situation of identical particles, *i.e.* fermions or bosons, the Hilbert space is restricted by the permutation (anti-) symmetry. Applying MCTDH directly³ to such a system is rather inefficient due to the redundancies in the expansion coefficients. However, it is possible to modify MCTDH by taking into account the permutation symmetries, which can significantly improve the efficiency. In the following section we will focus on the treatment of bosonic systems and briefly introduce the MCTDHB, which specializes MCTDH to the bosonic system for achieving higher performance.

B. MCTDHB

We now consider a single-species system of N bosons. All bosons share the same set of time-dependent orthonormal orbitals $\{|\phi_i(t)\rangle\}_{i \in [1,M]}$. In MCTDHB [23–25], the many-body wave function is then expanded within the basis of permanents, *i.e.* bosonic number states $|\vec{n}; t\rangle$, which naturally takes the bosonic symmetry into account and, thus, reduces the number of expansion coefficients in comparison to a MCDTH expansion. Omitting all

³ Since the Hamiltonian for a system of identical particles preserves the (anti-) symmetry of the total wave function during propagation, so does a converged MCTDH propagation.

time-dependencies in the notation as always, this yields the following ansatz wave function:

$$|\Psi\rangle = \sum_{\vec{n}|N} C_{\vec{n}} |\vec{n}\rangle, \quad (16)$$

where $\vec{n} = (n_1, \dots, n_M)$ and n_i is the occupation number of bosons in the orbital $|\phi_i\rangle$. The symbol “ $\vec{n}|N$ ” shall indicate that the summation runs over all positive integers n_i with the constraint $\sum_{i=1}^M n_i = N$. The number states $|\vec{n}\rangle$ become time-dependent as they equal permanents of time-dependent orbitals:

$$|\vec{n}\rangle = (N!n_1!\dots n_M!)^{-\frac{1}{2}} \sum_{\pi \in S(N)} |\phi_{i_{\pi(1)}}\rangle_1 \cdots |\phi_{i_{\pi(N)}}\rangle_N. \quad (17)$$

This equation is based on a multi-index $I = (i_1, \dots, i_N)$ which is fixed such that j occurs n_j times for any $j \in [1, M]$. The summation then runs over all permutations π of the first N integers, i.e. $\pi : [1, N] \rightarrow [1, N]$, and ket vectors get another index, i.e. $|\rangle_\kappa$, encoding to “which” boson the state belongs to. Furthermore, the time-dependent orbitals are in turn expanded with respect to some time-independent PBFs $\{|x\rangle\}$:

$$|\phi_i\rangle = \sum_x \phi_i(x) |x\rangle. \quad (18)$$

Next, we extend the diagrammatic representation of the total wave function to expansions in terms of permanents, which will turn out to be useful when discussing the expansion schemes of ML-MCTDHB: In the following, we indicate with a plus in a node that its SPFs⁴ are expanded in permanents. Then, the number of vertical lines below that bosonic node represents the number of bosonic particles, while the number of teeth of its child node equals the number of single particle orbitals. An example for a diagrammatic representation of a MCTDHB wave function for a system of six bosons in one-dimensional coordinate space is given in figure 2.

When restricting to at most two-body interactions, the many-body Hamiltonian can be expressed in the second-quantization picture as follows:

$$\hat{H} = \sum_{k,q} h_{kq} \hat{a}_k^\dagger \hat{a}_q + \frac{1}{2} \sum_{k,s,l,q} v_{kslq} \hat{a}_k^\dagger \hat{a}_s^\dagger \hat{a}_l \hat{a}_q \quad (19)$$

where $\hat{a}_k^\dagger(\hat{a}_k)$ is the time-dependent bosonic creation (annihilation) operator corresponding to the time-dependent orbital $|\phi_k\rangle$ and obeying $[\hat{a}_s, \hat{a}_q] = 0$, $[\hat{a}_s, \hat{a}_q^\dagger] = \delta_{sq}$. The time-dependent

⁴ Note that in a MCTDHB tree, the top node is always bosonic and represents a single SPF, namely the total wave function.




Figure 2: Visualization of a MCTDHB wave function expansion for a system of $N = 6$ bosons, which may occupy $m = 3$ orbitals. Each orbital is represented by $n = 5$ time-independent PBFs.

matrix elements of the one-body Hamiltonian $\hat{h}$ and two-body interaction potential $\hat{v}$, *i.e.* h_{kq} and v_{kslq} respectively, are defined as:

$$\begin{aligned} h_{kq} &= \langle \phi_k | \hat{h} | \phi_q \rangle, \\ v_{kslq} &= \langle \phi_k | \langle \phi_s | \hat{v} | \phi_l \rangle | \phi_q \rangle, \end{aligned} \quad (20)$$

and have to be evaluated in the representation (18).

The equations of motion for $C_{\vec{n}}$ and $\{|\phi_k(t)\rangle\}$ can be derived from the Dirac-Frenkel variational principle and read:

$$i\partial_t C_{\vec{n}} = \sum_{\vec{m}|N} \langle \vec{n} | \hat{H} | \vec{m} \rangle C_{\vec{m}}, \quad (21)$$

$$i\partial_t |\phi_k\rangle = (1 - \hat{P}) \left[\hat{h} |\phi_k\rangle + \sum_{j,s,l,q=1}^M [\rho^{-1}]_{kj} \rho_{jslq} \hat{v}_{sl} |\phi_q\rangle \right]. \quad (22)$$

In the latter equation, $\hat{P}$ is the projection operator onto the subspace spanned by the SPFs $\{|\phi_i\rangle\}$. ρ_{jk} denotes the element of the so-called one-body density matrix, defined as $\rho_{jk} = \langle \Psi | \hat{a}_j^\dagger \hat{a}_k | \Psi \rangle$, while ρ_{kslq} is the element of the so-called two-body density matrix, defined as $\rho_{kslq} = \langle \Psi | \hat{a}_k^\dagger \hat{a}_s^\dagger \hat{a}_l \hat{a}_q | \Psi \rangle$. Finally, the one-body Hamiltonian $\hat{h}$ as well as the mean field operator $\hat{v}_{sl}$ can be expressed in terms of the PBFs as:

$$\hat{h} = \sum_{x_1, x_2} \langle x_1 | \hat{h} | x_2 \rangle |x_1\rangle \langle x_2|, \quad (23)$$

$$\hat{v}_{sl} = \sum_{x_1, x_2} \langle x_1 | \langle \phi_s | \hat{v} | \phi_l \rangle | x_2 \rangle |x_1\rangle \langle x_2|. \quad (24)$$

By projecting (22) on a PBF $|x\rangle$, one arrives at the equation of motion for the coefficients $\phi_i(x)$. We note that (22) is equivalent to the ML-MCTDH non-top node equations of motion (12). The special form of (22) is a consequence of the Hamiltonian (19) containing only up to two body operators. The approach sketched above can be generalized to bose-bose mixtures and bose-fermi mixtures, and we refer the reader to [24, 25] for further details.

C. ML-MCTDHB

ML-MCTDH has proven to be a powerful tool for simulating complex systems with many degrees of freedom, while MCTDHB is optimized for treating scenarios involving one bosonic species. In this chapter, we derive the novel ML-MCTDHB method which both explicitly considers the bosonic symmetry of the particles and allows for a flexible adaption to system specific correlations and geometric aspects.

1. Concepts of ML-MCTDHB

While the wave function decomposition of ML-MCTDH is arbitrary in principle, ML-MCTDHB is based on two classes of wave function expansions: (i) species multi-layer and (ii) particle multi-layer MCTDHB. A general ML-MCTDHB wave function expansion then turns out to be a combination of these two classes. In the following, we discuss these classes before deriving the equations of motions:

(i) *Species multi-layer MCTDHB*: If the system under consideration consists of S bosonic species, $S > 1$, the total wave function is expanded in Hartree products $\prod_{\kappa=1}^S |\phi_{i_\kappa}^{2;\kappa}\rangle$:

$$|\Psi\rangle = \sum_{i_1=1}^{M_1} \dots \sum_{i_S=1}^{M_S} A_{i_1, \dots, i_S}^1 \prod_{\kappa=1}^S |\phi_{i_\kappa}^{2;\kappa}\rangle. \quad (25)$$

In the following, we refer to $|\phi_i^{2;\kappa}\rangle$ as the i -th species SPF of species κ . Hence, the expansion in terms of Hartree products reflects the distinguishability of the species. Each species SPF $|\phi_i^{2;\kappa}\rangle$ is in turn represented as a superposition of permanents $|\vec{n}\rangle$, paying respect to the bosonic symmetry:

$$|\phi_i^{2;\kappa}\rangle = \sum_{\vec{n}|N_\kappa} A_{i;\vec{n}}^{2;\kappa} |\vec{n}\rangle. \quad (26)$$

Here, $\vec{n} = (n_1, \dots, n_{m_\kappa})$ denotes the occupation numbers of the m_κ single particle orbitals, which can be populated by the bosons of species κ . The notation $\vec{n}|N_\kappa$ indicates the constraint that the occupation numbers n_i have to sum up to the number of bosons of species κ , i.e. N_κ . Finally, the SPFs underlying the time-dependent permanents $|\vec{n}\rangle$ are denoted by $|\phi_i^{3;\kappa}\rangle$, $i \in \{1, \dots, m_\kappa\}$, and expanded with respect to time-independent basis functions as usual for the SPFs of a primitive layer in ML-MCTDH. In the following, this class of wave function decomposition is denoted as *species multi-layer*.

The strength of species multi-layer MCTDHB lies in its flexibility with respect to the species SPFs: If the intra-species correlations are much stronger than the inter-species ones, only a few species many-body states, which are variationally optimally rotated by the ML-MCTDHB equations of motion, might be sufficient for a fair representation of the total state. So depending on the details of the system, species multi-layer MCTDHB allows to choose the optimal number of species SPFs, ranging from $M = 1$, the mean field limit for weakly interacting species, up to the full CI-limit for strongly correlated species when $M = \binom{N+m-1}{m-1}$.

A general species multi-layer expansion for a system of S species, each of them consisting of N_κ bosons, is shown in figure 3: For each species $\kappa \in \{1, \dots, S\}$, M_κ species SPFs are provided and the N_κ bosons constituting species κ may occupy m_κ orbitals. Although theses m_κ orbitals are expanded in n_κ time-independent states in figure 3, it should be pointed out that primitive mode combination can also be applied on this layer for a natural treatment of two- and three-dimensional bosons or internal degrees of freedom.

(ii) *Particle multi-layer MCTDHB*: A multi-layer expansion can also be applied to the orbitals of the bosons if these are higher-dimensional objects living in two- or three-dimensional coordinate space and/or having internal degrees of freedom. This multi-layer extension allows for grouping strongly correlated particle degrees of freedom together leading to a tailored representation of the bosonic orbitals for a given system.

Figure 4 shows some examples for particle multi-layer MCTDHB: The dashed lines heading to the bosonic nodes shall indicate that the shown tree diagrams may be incorporated in a species multi-layer MCTDHB expansion. Particularly, we would like to highlight how quasi-one or -two dimensional systems can be treated in quite a natural way: Whereas all three spatial degrees of freedom are treated equivalently in figure 4(a), there is an intermediate layer consisting of the combined xy and the z degrees of freedom in figures 4(c)




Figure 3: Archetype of a species multi-layer MCTDHB expansion of the wave function expansion for a system of S species, each of them consisting of N_κ bosons, which may occupy m_κ orbitals, which in turn are expanded in n_κ time-independent basis function. Each species may populate M_κ species SPFs. Please note that the vertical and horizontal lines are only schematic because the numbers of bosons, SPFs and PBFs are arbitrary. As S is arbitrary, the dashed lines shall indicate the expansion of the species κ' with

$$1 < \kappa' < \kappa \text{ or } \kappa < \kappa' < S.$$

and 4(d). This allows for associating much fewer states, here only a single one, to the energetically frozen out z degree of freedom while providing more states for xy degrees of freedom where correlated dynamics might be expected. The two-dimensional xy SPFs can then either be represented by primitive mode combination (figure 4(d)) or by a multi-layer expansion, which can be favourable for bosons in traps being asymmetric with respect to the xy plane (figure 4(c)). Furthermore, the particle multi-layer expansion easily allows for simulating bosons with internal degrees of freedom. Figure 4(b) illustrates this with an example of one-dimensional, unpolarized spin-one bosons. As the z component of the spin is not assumed to be fixed, the spin degree of freedom S may occupy as many states as given time-dependent basis functions, namely three.

After having introduced the species- and particle multi-layer MCTDHB wave function expansions, we briefly comment on two further tree diagram types: Naturally, the question




Figure 4: Examples for particle multi-layer MCTDHB: Figures a), c) and d) represent a species of bosons living in three-dimensional coordinate space. The wave function expansion a) reflects the equivalence of all three spatial dimensions, while in figures c) and d) the z -component is singled out by the fact that only one z orbital is provided. In figure c), the xy SPFs are expanded according to the ML-MCTDH philosophy treating both the x and the y degree of freedom equivalently. In figure d), mode combination is applied for expanding the xy SPFs. A particle multi-layer expansion is used for describing one-dimensional spin-one bosons in figure b). The spin degree of freedom is denoted by S . The dashed lines in all figures shall indicate that the shown species branch could be a part of a species multi-layer expansion.

arises whether tree diagrams are possible in which a bosonic node has bosonic children nodes for describing bosons made out of bosons. In the outlook, an extension of ML-MCTDHB to fermionic species will be discussed. Especially in this context, the question above gains importance as such trees could possibly mimic the forming of cooper pairs out

of electrons. Unfortunately, such tree structures break the permutation (anti-) symmetry of the constituting bosons (or fermions). Hence, this class of wave function expansion can only be applied as an approximation being valid under certain circumstances and will not be considered in this work.

Finally, we would like to point out that the second layer in species ML-MCTDHB, i.e. the species layer, does not need to consist only of bosonic nodes alone but can also contain normal nodes representing additional distinguishable degrees of freedom. In such a setting, we would say that the bosonic species nodes span bosonic subbranches in the tree presentation, while the normal nodes of the second layer induce normal subbranches. In the following section, it will become clear how one can easily derive the equations of motion for the coefficients belonging to nodes of a normal and a bosonic subbranch.

Summarizing, ML-MCTDHB makes use of the multi-layer concept above the bosonic nodes or below the child node of a bosonic node. Whereas the former can be used for an tailored treatment of inter-species correlation, the latter can be employed for reflecting the geometry of the system in coordinate space, i.e. the trap, and/or for considering internal degrees of freedom.

2. Correspondence principle between ML-MCTDH and ML-MCTDHB

The ML-MCTDHB equations of motion, which lead to a variationally optimal representation of the total wave function at each instant, can be derived by applying the Dirac-Frenkel variation principle to the ML-MCTDHB wave function ansatz. From a technical point of view, however, it is much easier to study the relationship between ML-MCTDHB and ML-MCTDH firstly and to deduce the ML-MCTDHB equations of motion from ML-MCTDH thereafter:

When dealing with systems containing indistinguishable bosons, a natural way of mode combination is to group all the bosons of the same species into one species node, *i.e.* one bosonic node, and this is the common starting point of ML-MCTDH and ML-MCTDHB. The difference between these two methods is that the SPFs of the species node are expanded with respect to permanents in ML-MCTDHB rather than Hartree products (see (26)), and consequently all identical bosons are represented by a single subnode below the species node. The correspondence principle now states the following: (i) Any ML-MCTDHB expansion can

be unravelled into a corresponding ML-MCTDH expansion. (ii) The expansion coefficients of non-bosonic nodes in ML-MCTDHB are the same as their counterparts in ML-MCTDH, and there is a linear mapping between the coefficients of a bosonic node in ML-MCTDHB and their ML-MCTDH counterparts. Consequently, we may translate the ingredients of the equations of motion of section IIA and the equations itself into the ML-MCTDHB framework.

To demonstrate the mapping between the bosonic coefficients and their counterparts, we consider a subbranch of a ML-MCTDH expansion, which contains N identical bosons. Paying respect to their indistinguishability, we provide m orbitals for each boson and group them on the same layer in the ML-MCTDH tree. Such a combination of all the bosonic DOF to a single logical DOF, the species DOF, appears quite natural as the permutation symmetry of the bosons enforces strong correlations between them.

Each species SPF⁵, say $|\psi_r\rangle$ for simplifying the notation, is expanded in terms of Hartree products made of the N SPFs. Each factor of such a Hartree product is an element of the set $\{|\varphi_i\rangle | i \in [1, m]\}$. The indistinguishability of the bosons is then reflected in the symmetry of the expansion coefficients $A_{r;I}$ with respect to permutation within the multi-index $I = (i_1, \dots, i_N)$. As a consequence, the SPF $|\psi_r\rangle$ can be equivalently expanded in permanents made of the orbitals $|\varphi_i\rangle$:

$$|\psi_r\rangle = \sum_I A_{r;I} |\varphi_{i_1}\rangle \cdots |\varphi_{i_N}\rangle = \sum_{\vec{n}|N} \tilde{A}_{r;\vec{n}} |\vec{n}\rangle. \quad (27)$$

This identity implies that for each such a ML-MCTDH expansion of a bosonic species there is an equivalent ML-MCTDHB expansion and vice versa. An example for this correspondence principle is given in figure 5. Since we require the permanents to be orthonormal, i.e. $\langle \vec{m} | \vec{n} \rangle = \delta_{\vec{m}, \vec{n}}$, we arrive at the following correspondence relation between the ML-MCTDH and ML-MCTDHB coefficients $A_{r;I}$ and $\tilde{A}_{r;\vec{n}}$, respectively:

$$\tilde{A}_{r;\vec{n}} = \sqrt{\frac{N!}{\prod_{i=1}^m n_i!}} A_{r;I}, \quad (28)$$

where I is some multi-index, which contains n_s times the number s , $s \in [1, m]$. This translation rule ensures the orthonormality of the species SPFs in both representations, i.e. $\sum_I A_{k;I}^* A_{r;I} = \delta_{kr}$ if and only if $\sum_{\vec{n}} \tilde{A}_{k;\vec{n}}^* \tilde{A}_{r;\vec{n}} = \delta_{kr}$. Because of this one-to-one correspondence between the coefficients, we omit the tilde in the notation. The translation of the

⁵ Or the total wave function in case of only a single species.




Figure 5: Visualization of the correspondence principle between ML-MCTDHB and ML-MCTDH: In subfigure a), a bosonic branch of a ML-MCTDHB wave function expansion is shown. This branch represents three bosons, which may occupy two orbitals. The dotted lines starting from the bosonic node as well as from its child node shall indicate that the shown branch may be incorporated in a species- or particle multi-layer MCTDHB expansion or both. Subfigure b) shows the corresponding ML-MCTDH representation of this branch. Please note that each of the three children nodes is associated with two SPFs because of the indistinguishability of these DOF. Due to the same argument, the very same subtree must be beneath each of these children nodes, which is again indicated by the two dotted lines.

equations of motion from ML-MCTDH to ML-MCTDHB mainly subjects to the replacement of the summation over Hartree products of the species node to that over permanents. The replacement can be illustrated as:

$$\sum_I f(A_{i;I}) = \sum_{\vec{n}} \frac{N!}{\prod_{i=1}^m n_i!} f\left(\sqrt{\frac{\prod_{i=1}^m n_i!}{N!}} A_{r;\vec{n}}\right), \quad (29)$$

where f is an arbitrary function. Following this, we can see that the equations of motion for the DOF above and below the bosonic node(s) are identical between ML-MCTDH and ML-MCTDHB. So the major difference between the two methods shows up in the bosonic node(s), where the density and the mean field (operator) matrices have to be expressed in the second quantization language as well as their action on the SPFs. In the following sections, we will apply the correspondence principle in order to derive the equations of motion for species- and particle ML-MCTDHB.

Before we step to the concrete equations of motion of ML-MCTDHB, we would like to

comment briefly on conservation laws: As shown in the appendix of [8], all sets of equations of motion which can be derived from the Dirac-Frenkel variational principle preserve both the norm and the energy of the total wave function. In addition to this, we have proven analytically that the equations of motion of the MCTDH family preserve a certain class of symmetries in the following sense: Let the Hamiltonian $\hat{H}$ have some symmetry group $\mathcal{G}$ and let $\hat{G}$ be an element of the unitary representation of $\mathcal{G}$ in the many-body Hilbert space, i.e. $[\hat{G}, \hat{H}] = 0$. Here, we only consider symmetry operations which can be decomposed as $\hat{G} = \bigotimes_i \hat{g}_i$, where $\hat{g}_i$ is an element of the unitary representation of $\mathcal{G}$ in the Hilbert space of the i -th physical DOF⁶. For simplicity, we focus on the MCTDH method⁷. If at $t = 0$ the many-body wave function $|\Psi(0)\rangle$ and all the SPFs $|\phi_j(0)\rangle$ are now invariant under $\hat{G}$ and $\hat{g}_j$, respectively, they remain invariant forever⁸, i.e. $\hat{G}|\Psi(t)\rangle = e^{i\Theta}|\Psi(t)\rangle$ and $\hat{g}_j|\phi_j(t)\rangle = e^{i\theta_j}|\phi_j(t)\rangle$ for all t . This property of the MCTDH family cannot be expected *a priori* due to the truncation of the many-body Hilbert space and the complicated integro-differential equations for the SPFs and we are not aware of any work which has shown this before. Possibly this symmetry conservation can be employed for specifying the configuration selection in selected configuration MCTDH type methods (cf. [38] and references therein). For a sketch of the proof, see appendix A.

3. Species Multi-Layer MCTDHB

In this section we present the theory of multi-layer MCTDHB in detail, and firstly we focus on *species multi-layer*, considering a mixture system of S species confined in one-dimension, as shown in figure 2. As indicated in equation (25), the total wave function is expanded as:

$$|\Psi\rangle = \sum_{i_1=1}^{M_1} \dots \sum_{i_S=1}^{M_S} A_{i_1, \dots, i_S}^1 \prod_{\kappa=1}^S |\phi_{i_\kappa}^{2;\kappa}\rangle \equiv \sum_{I=(i_1, i_2, \dots, i_S)} A_I^1 |I^1\rangle, \quad (30)$$

where all time-dependencies are omitted as always. The species SPF $|\phi_i^{2;\kappa}\rangle$ is in turn represented in terms of permanents $|\vec{n}\rangle_\kappa$ defined as in equation (17) :

$$|\phi_i^{2;\kappa}\rangle = \sum_{\vec{n}|N_\kappa} A_{i;\vec{n}}^{2;\kappa} |\vec{n}\rangle_\kappa. \quad (31)$$

⁶ Some of the $\hat{g}_i$ -th may be the unit operator.

⁷ The sketched proof also holds for MCTDHB, MCTDHF and ML-MCTDH and thus also for ML-MCTDHB.

⁸ In ML-MCTDH, the invariance of the SPFs on all layers under the corresponding symmetry operations has to be assumed at $t = 0$ and will then be preserved forever.

The index κ of the permanent shall indicate that $|\vec{n}\rangle_\kappa$ is a permanent made of the SPFs $|\phi_i^{3;\kappa}\rangle$ which are associated with the physical DOF of species κ bosons, i.e. x_κ . In the end, these SPFs $|\phi_i^{3;\kappa}\rangle$ are expanded with respect to the time-independent PBFs $|x_r^\kappa\rangle$:

$$|\phi_i^{3;\kappa}\rangle = \sum_r A_{i,r}^{3;\kappa} |x_r^\kappa\rangle. \quad (32)$$

Without loss of generality, we restrict ourselves to Hamiltonians involving at most two-body terms - a generalization to $R > 2$ body interactions is straightforward. So we assume a Hamiltonian of the form:

$$\hat{H} = \sum_{\kappa=1}^S [\hat{H}_\kappa + \hat{V}_\kappa] + \sum_{\kappa < \kappa'} \hat{W}_{\kappa,\kappa'}, \quad (33)$$

where $\hat{H}_\kappa$ contains all one-body terms of species κ , i.e. $\hat{H}_\kappa = \sum_{i=1}^{N_\kappa} \hat{h}_i^\kappa$ with $\hat{h}_i^\kappa$ acting only on the i -th κ boson. $\hat{V}_\kappa$ contains all intra-species interaction terms of species κ and equals $\sum_{i < j} \hat{v}_{ij}^\kappa$ with $\hat{v}_{ij}^\kappa$ affecting only the i -th and j -th κ boson. Finally, $\hat{W}_{\kappa,\kappa'}$ denotes the inter-species interaction between the species κ and κ' and equals $\sum_{i=1}^{N_\kappa} \sum_{j=1}^{N_{\kappa'}} \hat{w}_{i,j}^{\kappa,\kappa'}$ with $\hat{w}_{i,j}^{\kappa,\kappa'}$ acting only on the i -th κ and the j -th κ' boson. Without loss of generality, all two-body operators are assumed to be given as a sum over products of two one-body operators:

$$\begin{aligned} \hat{v}_{ij}^\kappa &= \sum_\nu C_\nu^\kappa \hat{v}_i^{\kappa,\nu} \hat{v}_j^{\kappa,\nu}, \\ \hat{w}_{i,j}^{\kappa,\kappa'} &= \sum_\nu D_\nu^{\kappa,\kappa'} \hat{w}_i^{\kappa,\nu} \hat{w}_j^{\kappa',\nu}. \end{aligned} \quad (34)$$

Such a form is particularly handy for avoiding high-dimensional integrals when calculating matrix elements and for the sake of a recursive formulation of the ML-MCTDHB ingredients.

For applying the correspondence principle, we firstly summarize that the total wave function and each SPF is expressed as a summation over coefficients times states $|C\rangle$, which are either Hartree products (for the top layer), permanents (for the species layer) or PBFs (for the physical layer). According to the correspondence principle, the top and the non-top node equations of motion, respectively, then read:

$$i\partial_t A_C^1 = \sum_D \langle C | \hat{H} | D \rangle A_D^1, \quad (35)$$

$$i\partial_t A_{n;C}^{\alpha;\kappa} = \sum_D \langle C | (1 - \hat{P}^{\alpha;\kappa}) \sum_m [(\rho^{\alpha;\kappa})^{-1} * \langle \hat{H} \rangle^{\alpha;\kappa}]_{n,m} A_{m;D}^{\alpha;\kappa} | D \rangle, \quad (36)$$

where the density matrix, the mean field operator matrix and the projector are given by $\rho_{n,m}^{\alpha;\kappa} = \langle \Psi_n^{\alpha;\kappa} | \Psi_m^{\alpha;\kappa} \rangle$, $\langle \hat{H} \rangle_{n,m}^{\alpha;\kappa} = \langle \Psi_n^{\alpha;\kappa} | \hat{H} | \Psi_m^{\alpha;\kappa} \rangle$ and $\hat{P}^{\alpha;\kappa} = \sum_i \sum_{E,F} (A_{i;F}^{\alpha;\kappa})^* A_{i;E}^{\alpha;\kappa} | E \rangle \langle F |$, respectively. Here, the SHFs are defined in the same way as in (9) everywhere in the tree but

for bosonic particle nodes, where we have the definition⁹: $|\Psi_m^{\alpha;\kappa}\rangle := \frac{\hat{a}_{\kappa,i}}{\sqrt{N_\kappa}}|\Psi\rangle$. Consequently, we are left to evaluate those ingredients as well as the top layer mean field $\langle C|\hat{H}|D\rangle$. In appendix B, we give in detail the equations of motion for the coefficients of different layers as well as the formulas for all the relevant ingredients.

4. Particle Multi-Layer MCTDHB

In this section we turn to particle ML-MCTDHB, which is mainly designed for bosonic systems with a high-dimensional configuration space, spanned by the spatial DOF, and/or internal DOF. To demonstrate the implementation of the particle-ML MCTDHB, we consider a single-species bosonic system of N identical bosons living in a Q -dimension configuration space, spanned by the physical DOF $\{x_1, x_2, \dots, x_Q\}$. In ML-MCTDHB, we firstly perform a mode combination of these physical DOF, and generate the logical DOF one layer above the physical layer, as in equation (2), and repeat the mode combination recursively layer by layer, until the layer containing only one node, which refers to the DOF of the boson $\vec{x} = (x_1, x_2, \dots, x_Q)$. Above this particle node, there is the top node representing the state vector $|\Psi\rangle$ of the whole system. This mode-combination procedure corresponds to the tree structure shown in figure 6, where the dashed box between the particle node and the physical nodes indicates the variety of possible mode-combination schemes. As an example, we have shown different mode-combination schemes of two or three physical DOF into one particle DOF in figure 4.

We associate a set of SPFs to each logical DOF on the non-top layer, and expand them with respect to the SPFs of the corresponding DOF one layer below, as:

$$|\phi_n^{\alpha;\kappa}\rangle = \sum_{i_1, i_2, \dots, i_m} A_{n; i_1, i_2, \dots, i_m}^{\alpha;\kappa} \prod_{r=1}^m |\phi_{i_r}^{\alpha+1;\kappa_r}\rangle \equiv \sum_I A_{n; I}^{\alpha;\kappa} |I^{\alpha;\kappa}\rangle, \quad (37)$$

where $|\phi_n^{\alpha;\kappa}\rangle$ is the n -th SPF of the κ -th DOF on the α -th layer. The state vector $|\Psi\rangle$ of the top node is, however, expanded with respect to permanents :

$$|\Psi\rangle = \sum_{\vec{n}} A_{\vec{n}}^1 |\vec{n}\rangle, \quad (38)$$

⁹ Please note that although the SHF for a bosonic node is given in the second quantization representation, we exclusively treat problems with fixed particle numbers in the ML-MCTDHB formalism. This in turn means that projections like $\langle \Psi_i^{3;\kappa} | \Psi \rangle$ have to be understood as an incomplete scalar product in the $\sum_{\sigma} N_{\sigma}$ particle Hilbert space resulting in a one-particle state and not as the scalar product in the Fock space.




Figure 6: Illustration of a general particle multi-layer MCTDHB wave function expansion for spin-one bosons in three-dimensional coordinate space. The dashed box indicates the various possibilities for mode combination of the physical DOF, i.e. the spatial DOF x, y, z and the spin projection on the z -axis.

where $|\vec{n}\rangle$ is the number state as defined in equation (17). The physical DOF are also associated with a set of SPFs, and these SPFs are expanded with respect to time-independent PBFs as in ML-MCTDH (see equation (5)). If the physical DOF x_κ is a scalar quantity like the position in one direction, the associated SPFs are represented by one-dimensional PBFs. In the case of strong correlations treated with primitive mode combination, i.e. if the physical DOF x_κ is vector-valued, its SPFs are still expanded with respect to PBFs, but each PBF is then a product of one-dimensional time-independent basis vectors:

$$|\phi_j^{P;\kappa}\rangle = \sum_{x_\kappa} A_{j;x_\kappa}^{P;\kappa} |x_\kappa\rangle \equiv \sum_{I=(i_1,\dots,i_n)} A_{j;I}^{P;\kappa} \prod_{\sigma=1}^n |x_{i_\sigma}^{\kappa,\sigma}\rangle, \quad (39)$$

An example for incorporating primitive mode combination in particle multi-layer MCTDHB is shown in figure (4(d)), where the x and y DOF share the same set of SPFs, and are grouped into one primitive node.

For simplicity, we restrict ourselves to two-body interactions and, moreover, assume the

product representation of the Hamiltonian as usually:

$$\begin{aligned}\hat{H} &= \sum_i \hat{h}(i) + \sum_{i<j} \hat{V}(i, j) \\ &= \sum_i \hat{h}(i) + \sum_{i<j} \sum_{\nu} C_{\nu} \hat{v}_{\nu}^1(i) \hat{v}_{\nu}^2(j).\end{aligned}\tag{40}$$

$\hat{h}(i)$ and $\hat{V}(i, j)$ are the one-body Hamiltonian acting on the i -th boson, and the two-particle interaction operator acting on the i -th and j -th bosons respectively, where the two-particle interaction operator is written as the summation of products of single-particle operators acting on the i -th and j -th boson separately. The single-particle operators $\hat{h}$, $\hat{v}_{\nu}^i$ ($i=1,2$) can be further expressed as the summation of products of operators acting on each of the physical DOF, e.g.:

$$\hat{h} = \sum_{\mu} D_{\mu} \prod_{\kappa} \hat{h}_{\mu}^{\kappa},\tag{41}$$

where $\hat{h}_{\mu}^{\kappa}$ is a operator acting on the physical DOF x_{κ} .

According to the correspondence principle, the equations of motion for the non-top nodes are equivalent to those in ML-MCTDH, and the top-node equations of motion are expressed in the second quantization picture. The equations of motion of all the coefficients can be given as following:

$$\begin{aligned}i\partial_t A_{\vec{n}}^1 &= \sum_{\vec{m}} \langle \vec{n} | \hat{H} | \vec{m} \rangle A_{\vec{m}}^1, \\ i\partial_t A_{n;I}^{\alpha;\kappa} &= \sum_J \langle I^{\alpha;\kappa} | (1 - \hat{P}^{\alpha;\kappa}) \hat{h}_{\nu}^{\alpha;\kappa} | J^{\alpha;\kappa} \rangle A_{n;J}^{\alpha;\kappa} \\ &\quad + \sum_{\nu} C_{\nu} \{ \sum_J \langle I^{\alpha;\kappa} | (1 - \hat{P}^{\alpha;\kappa}) \hat{v}_{\nu;1}^{\alpha;\kappa} | J^{\alpha;\kappa} \rangle \} \\ &\quad \times \{ \sum_m [(\rho^{\alpha;\kappa})^{-1} * \langle \hat{v}_{\nu;2}^{\alpha;\kappa} \rangle]_{n,m} \} A_{m;J}^{\alpha;\kappa}.\end{aligned}\tag{42}$$

For the equations of motion of the non-top node, $\hat{P}^{\alpha;\kappa} \equiv \sum_n |\phi_n^{\alpha;\kappa}\rangle \langle \phi_n^{\alpha;\kappa}|$ is the projection operator, and $\hat{h}^{\alpha;\kappa}$ and $\hat{v}_{\nu;i}^{\alpha;\kappa}$ ($i=1,2$) are the single-particle operators of the one-body Hamiltonian and two-particle interactions, which are defined recursively as shown in the ML-MCTDH section (cf. equation (8)). $\rho^{\alpha;\kappa}$ and $\langle \hat{v}_{\nu;2}^{\alpha;\kappa} \rangle$ denote the density and mean field matrices of the corresponding node, respectively. For the particle node, the density matrix is calculated as:

$$\rho_{ij}^{2;1} = \frac{1}{N} \sum_{\vec{n}|N-1} \sqrt{(n_i + 1)(n_j + 1)} (A_{\vec{n}+\hat{i}}^1)^* A_{\vec{n}+\hat{j}}^1,\tag{43}$$

and the mean field matrices for the two-particle interaction terms are calculated as:

$$\begin{aligned} \langle \hat{v}_{\nu;2}^{2;1} \rangle_{i,j} = & \frac{1}{N} \sum_{\vec{n}|N-2} \sum_{p,q} \sqrt{(n_i+1)(n_p+\delta_{i,p}+1)} \sqrt{(n_j+1)(n_q+\delta_{j,q}+1)} \\ & * (A_{\vec{n}+\hat{i}+\hat{p}}^1)^* A_{\vec{n}+\hat{j}+\hat{q}}^1 \langle \phi_p^{2;1} | \hat{v}_{\nu}^2 | \phi_q^{2;1} \rangle. \end{aligned} \quad (44)$$

The density and mean field matrices of the nodes under the particle node are generated as in ML-MCTDH (see equations (13,14)), which is again related to the correspondence principle and enables one to apply the well-known recursion schemes [15, 16].

By now we have introduced the particle-ML MCTDHB with an example of a single-species bosonic system. The generalization to bosonic multispecies systems is straightforward, and we just need to apply the correspondence principle to the layers below the particle nodes, as we have done here for the single-species case, while we apply the species multi-layer to the layers above. This way we also complete the merging of the species multi-layer MCTDHB with particle multi-layer MCTDHB, obtaining the ability to deal with bosonic systems of arbitrary species in arbitrary high dimensions.

As a final remark, we would like to point out the relationship between ML-MCTDHB and the standard MCTDHB. For single-species system, ML-MCTDHB and MCTDHB coincide for one-dimensional bosons without internal degrees of freedom. When considering bosons living in a higher-dimensional configuration space, ML-MCTDHB reduces to MCTDHB if primitive mode combination is applied directly below the particle node. For mixtures, the state vector of the total system is expanded in standard MCTDHB as the summation of the products of the permanents states belonging to each species, while in species multi-layer MCTDHB the state vector is firstly expanded with respect to Hartree products of species SPFs, and the species SPFs are then expanded within the permanents basis of each species. Hence, ML-MCTDHB recovers the standard MCTDHB treatment if as many species SPFs are provided as there are configurations for each species.

III. APPLICATIONS

Having introduced the theory of ML-MCTDHB, we now turn to some applications of the method. Our ML-MCTDHB code is based on the general ML-MCTDH implementation [16] in its recursive formulation [15]. Since this ML-MCTDH implementation is capable of handling arbitrary ML-MCTDH wave function expansions, we have extended the code to

species and particle multi-layer MCTDHB as well as mixtures of both. Hence, in principle an arbitrary number of bosonic species in arbitrary dimensions can be simulated as long as the necessary number of configurations is not too large.

At the moment, our ML-MCTDHB code can deal with arbitrary two-body interactions - an extension to three- or four-body interactions would be straightforward. As explained above, the recursive ML-MCTDH formulation demands a certain product representation of the interaction potential and so does our ML-MCTDHB implementation. This product representation can be achieved by the POTFIT algorithm [39, 40]. Since most of our further applications are situated in the field of ultra-cold bosonic atoms, the contact interaction will be extremely relevant. Approximating the delta interaction potential by a narrow Gaussian, the POTFIT machinery can be applied for obtaining the desired product form (cf. [12]). In two- and three-dimensional systems, a word of caution is in order here: For instance, when expressing the regularized contact interaction in three dimensions, $V(\vec{r} = \vec{x}_1 - \vec{x}_2) = g \delta^{(3)}(\vec{r}) \partial_r r$, in terms of the particle coordinates $\vec{x}_1, \vec{x}_2$, one also has to bring additional terms in the product form due to the modulus of $\vec{r}$ and its derivative with respect to the different components of $\vec{x}_1$ and $\vec{x}_2$. For one-dimensional settings, we have also developed an exact implementation of the contact interaction $V(x_1 - x_2) = g \delta(x_1 - x_2)$ *without* reshaping it into the product form in order to liberate the simulations from the artificial length scale induced by the Gaussian approximation of the delta function (cf. appendix C). The main difference in the derivation of the equations of motion lies in the fact that for a Hamiltonian being not of the product form (6) one cannot separate the mean field operators into a scalar mean field matrix and a operator valued factor anymore, which breaks the recursive multi-layer formulation.

Depending on the concrete problem, we employ an appropriate discrete variable representation (DVR) of the SPFs and the Hamiltonian [45]. So in the case of the exact implementation of the delta interaction, the grid point spacing defines the smallest length scale and, thus, an ultraviolet cutoff.

For integrating the equations of motion, we employ either ZVODE, a variable-coefficient ordinary differential equation solver with fixed-leading-coefficient implementation [46], or DOPRI, an explicit Runge-Kutta method of order five with stepsize control and dense output [47].

In the following, we focus on the tunneling dynamics in a one-dimensional double well

system made of a harmonic trap superimposed with a Gaussian at the trap centre, i.e. $V_{trap}(x) = x^2/2 + h/\sqrt{2\pi s^2} \exp(-x^2/2s^2)$ in harmonic oscillator units $\hbar = m = \omega = 1$. Firstly, we will show the correctness of our implementation by comparing the single component tunneling results with results from the well tested implementation of MCTDH [48]. Then we apply species ML-MCTDHB to simulate the tunneling of a binary mixture and compare with the established ML-MCTDH approach [16]. Afterwards, we turn to a more complex scenario of three species with higher number of particles studying signatures of correlations and discussing the scaling in comparison to MCTDH, ML-MCTDH and MCTDHB. Applications of particle ML-MCTDHB will be presented elsewhere.

In all examples, we prepare the initial state by modifying the trapping potential such that it becomes energetically favourable for the bosons to be in the left well. All bosons are then put into the ground state of the resulting single particle Hamiltonian and, afterwards, we let the many-body system relax to its ground state by propagating the equations of motion in imaginary time. The resulting many-body state is finally propagated in real time in the original double well trap.

A. Single-Species Tunneling

As a first application, we simulate the tunneling dynamics of a single-species bosonic ensemble in the double well. The tunneling dynamics of a bosonic ensemble in a one-dimensional double well has been extensively studied, for both microscopic and macroscopic systems, with boson numbers ranging from two to the order of 10^6 and even more [12, 31, 49–52]. Experiments on the double well tunneling have also been carried out [53–55]. Various theoretical approaches have been employed, for instance, the Gross-Pitaevskii equation [49, 50], the Bose-Hubbard model [51, 52] as well as ab-initio methods: MCTDHB has been applied to the many-body system [31], and calculations via MCTDH have been carried out for few-body systems [12]. The simulations based on the single-band approximation predict that the tunneling is suppressed for interaction strengths above some critical value, while the extended model predicts the weakening of such a suppression in the strong interaction regime, where higher bands effects cannot be neglected. In this way the double well potential manifests itself as a proper test bed for the higher band effects. In this section we will simulate the double well tunneling with ML-MCTDHB, and perform a detailed analysis to




Figure 7: The population oscillation of four bosons in the left and right well, with (a) $g=0.0$, (b) $g=0.5$, (c) $g=2.0$, and (d) $g=4.0$. For interaction strengths $g \leq 2.0$ (figures (a)-(c)), six orbitals are enough to achieve a good convergence, while in the case of $g = 4.0$ (figure (d)), ten orbitals are needed to meet the convergence requirement. The results of ML-MCTDHB are shown as solid lines, and the MCTDH results as crosses. A good agreement is observed between the results of both methods.

resolve the higher bands effects.

Here we present the tunneling dynamics of four and ten bosons in a double well potential. A sin-DVR is employed, which intrinsically introduces hard-wall boundaries at both ends of the potential (cf. appendix of [8]). Firstly we show the simulation of the four-boson tunneling, in comparison with the MCTDH simulation using the Heidelberg MCTDH package [48]. To have a direct comparison between ML-MCTDHB and MCTDH results, we adopt in both simulations a narrow Gaussian interaction to model the contact interaction, and the interaction is written as $V(x_1 - x_2) = g(2\pi\sigma^2)^{-1/2} \exp(-(x_1 - x_2)^2/(2\sigma^2))$, with $\sigma = 0.05$. Next we extend the simulation to the ten-boson case, and the contact interaction is modeled by the exact delta function. The simulation of the tunneling of ten bosons becomes impractical with MCTDH, and we only perform the simulation with ML-MCTDHB.

Figure 7 summarizes the population evolution of 4 bosons in the double well at different interaction strengths. When the interaction strength is zero, the system undergoes Rabi

oscillations, as shown in figure 7(a). As the interaction strength increases to 0.5, the oscillations almost vanish on a relative long time scale, which indicates the delayed tunneling behaviour. When the interaction strength increases to 2.0, as shown in figure 7(c), the amplitude of the population oscillations is increased from less than 0.06 in delayed tunneling (figure 7(b)) to around 1.0, which is referred to as enhanced tunneling. As the interaction increases even further to 4.0, the equilibration state emerges during tunneling, where the populations of the left and the right well approach the value of two with only small fluctuations, as shown in figure 7(d). Summarizing, figure 7 illustrates the tunneling transition from Rabi oscillations through delayed tunneling and enhanced tunneling to the equilibration state as the interaction strength increases from zero to the strong interaction regime. The ML-MCTDHB results show a very good agreement with the MCTDH calculations. Only for the interaction strength $g = 4.0$, deviations occur, which can be explained by the different implementations of the POTFIT algorithm in MCTDH and ML-MCTDHB. Nevertheless, we still observe qualitatively the same behaviour of the emergence of the equilibration state in both simulations. Moreover, more orbitals are needed to achieve good convergence in the strong interaction case of $g = 4.0$, and we supply ten orbitals to this case, where good convergence can be deduced from the natural populations discussed in the following.

The ML-MCTDHB simulation supplies the dynamical information of the system in terms of the time propagation of the wave function, and we can extract various dynamical properties of the system, via different analysis procedures. Among other things, we can calculate the eigenvalues and eigenvectors of the reduced one-body density matrix, the so-called natural populations and natural orbitals, respectively, as well as the one-body density profiles at different times during the tunneling. The natural populations can confirm the convergence of the calculation and also manifest themselves as a measure of the fragmentation of the system. To uncover the physical effect giving rise to the enhanced tunneling and equilibration state, we plot the natural populations and one-body density profiles of the four-boson ensemble at different time instants during the tunneling process in figures 7. Figure 8(a) and (c) show the natural populations and one-body densities for $g = 2.0$. In figure 8(a), firstly we see the lowest natural population saturates to a value less than 10^{-3} , which confirms the convergence of the simulation, and also that only two natural orbitals are macroscopically occupied. In figure 8(c), we observe that the profile in the left well remains as a Gaussian packet, while the profile in the right well presents a two-hump structure. The two-hump




Figure 8: The natural populations and one-body densities of the system of four bosons in the double well at different time instants, with (a) and (c) $g=2.0$, as well as (b) and (d) $g=4.0$.

profile is a signature of the occupation of the first excited state in the right well, and this indicates that the enhanced tunneling is due to the higher band occupation, *i.e.* the interband tunneling. In the 2-boson ensemble [12] the enhanced tunneling only takes place in the fermionization regime, *i.e.* for the interaction strength approaching infinity, while as the number of bosons increases, it becomes easier to excite higher bands, and the enhanced tunneling arises even in an interaction regime far below the fermionization limit.

The natural populations of $g = 4.0$, as shown in figure 8(b) shows a good convergence of the simulation with the lowest natural population saturating well below 1 percent, and at $g = 4.0$ more natural orbitals contribute to the tunneling process, which suggests that fragmentation of the system and decoherence occur during tunneling. Figure 8(d) shows the one-body densities at different times for the interaction strength $g = 4.0$, where the equilibration state dominates the tunneling. During the tunneling, the one-body density profile presents multiple wiggles in both the left and right wells, which indicates multiple higher-band excitations in the tunneling process. In the exact quantum dynamics study of the double well system [31], the equilibration state is explained by the decoherence of bosons between the left and right well, and the multi-excitation of higher energetic levels in the left




Figure 9: The population oscillation of ten bosons in the double well, with (a) $g=0.5$, (b) $g=1.0$. Enhanced tunneling is obtained in (a), where the amplitude of the population is around 2. The slow evolution to the equilibration state is obtained in (b).

and right well suggests that the tunneling process involves a large number of higher band number states, and in consequence multiple tunneling channels. The dephasing between these tunneling channels can be a source of the decoherence of the bosons in the two wells.

Figure 9 shows the tunneling evolution of the system of ten bosons in the double well with respect to the interaction strength. In the case of ten bosons, we focus on the enhanced tunneling and equilibration state for a sufficiently high interaction strength. The enhanced tunneling is observed with the interaction strength as weak as $g = 0.5$, as shown in figure 9, and at $g = 1.0$ the system slowly evolves to the equilibration state. Figures 10(a) to (d) show the natural populations and one-body densities at different time instants during the tunneling process of figures 9(a) and (b), respectively. The convergence of the simulations is demonstrated by the lowest natural population, which saturates to the order of 0.1 percent and 1 percent in figures 10 (a) and (b), respectively. In figure 10(c), the two-hump profile in the right well indicates that the enhanced tunneling is again as a result of interband tunneling, and the multi-wiggle structure in figure 10(d) suggests that multiple tunneling channels are involved and this can lead to the decoherence between the two wells, and consequently to the appearance of the equilibration state.




Figure 10: The natural population and one-body densities of ten bosons in the double well at different time, with (a) and (c) $g=2.0$, as well as (b) and (d) $g=4.0$.

B. Mixture Tunneling

Let us now study the tunneling dynamics of two and three bosonic species, which are loaded in the left well of the double well trap. All species shall have the same mass, which is set to one, and shall experience the same double well potential $V_{trap}(x)$. The intra-species interaction strengths g_σ of the contact interaction potentials $V_\sigma(x) = g_\sigma\delta(x)$, however, are assumed to be different for different species σ . Furthermore, the inter-species interaction is also modelled by a pseudo-potential: $V_{\sigma\sigma'}(x_\sigma - x_{\sigma'}) = g_{\sigma\sigma'}\delta(x_\sigma - x_{\sigma'})$. All delta potentials are implemented numerically exactly as sketched above. We choose $h = 3$ and $s = 0.2$ for the height and width of the barrier, respectively, which leads to three bands each consisting of two single particle eigenstates. The lowest band is separated by an energy difference of 1.63 from the first excited band and its level spacing amounts to $\Delta E \approx 0.23$ leading to a tunneling period of $T = 2\pi/\Delta E \approx 27$ for non-interacting particles. Assuming an equal number of particles of each species, $N_\sigma \equiv N$, and not too different intra-species interaction strengths, we provide for each species the same number of species SPFs, $M_\sigma \equiv M$, and particle SPFs, $m_\sigma \equiv m$. For preparing the initial state of the mixture, we modify the double well trap by letting $V_{trap}(x) = 20.0$ for $x > 0$ and also close to the left end point of the

spatial grid.

In order to illustrate the correctness of our implementation, we first consider a binary mixture made of 2 A and 2 B bosons. Here, we give an example where we have chosen $g_A = 0.3$, $g_B = 0.5 g_A$ and $g_{AB} = 0.1 g_A$. Furthermore, we allow each particle to occupy $m = 3$ particle SPFs and each species to be in $M = 4$ species states. In the following, we compare simulations of this system done with ML-MCTDHB and ML-MCTDH [16]. Although the initial state for the relaxation run is - from a mathematical perspective - perfectly symmetric with respect to particle exchange within each species, one has to pay attention to the initially unoccupied species SPFs in ML-MCTDH. These have to be symmetrized 'by hand' because otherwise the ML-MCTDH propagation will not preserve the exchange symmetry within each species.

MCTDH and its derivative methods are proven to preserve both the norm and the total energy exactly (cf. appendix of [8]). Using the ZVODE integrator with an absolute and relative tolerance of 10^{-10} and integrating 100 harmonic oscillator time units, the norm of the total wave function deviates from unity by a factor of 10^{-7} and the total energy is conserved up to a factor 10^{-6} for both ML-MCTDH and ML-MCTDHB.

Before discussing the convergence of the simulations with respect to the number of particle and species SPFs, we first summarize the results for different one- and two-body observables. Figure 11 shows the time evolution of the probability for finding an A (B) boson in the left well. One clearly sees that the tunneling period of the A bosons is enlarged in comparison to the Rabi-tunneling period, a manifestation of the well known delayed tunneling (cf. [12, 56]). In contrast to this, the probability evolution of finding a B boson in the left well qualitatively resembles a first Rabi-cycle but afterwards also features delayed tunneling. This observation is quite plausible: Since both the g_B and g_{AB} are relatively small in comparison to g_A , one expects that the B bosons require a longer interaction time in order to show an interaction induced effect. The impact of the different interaction strengths can also be seen in figure 12: While the probability for finding two A bosons in the same well is well above 0.5 for most of the propagation time, the B bosons tend to stay in the same well less likely. This bunching tendency of the A bosons might be interpreted as the onset of correlated pair tunneling induced by the intra-species interaction (cf. [12]). To the contrary, the probability for finding an A and a B boson in the same well fluctuates around 0.5 indicating that the bosons of each species tunnel independently.




Figure 11: 2 A and 2 B bosons initially loaded in the left well of a double well trap: The blue (red) line shows the probability for finding an A (B) boson in the left well versus time. Lines correspond to ML-MCTDHB results while crosses are obtained via ML-MCTDH calculations. Parameters are: $g_A = 0.3$, $g_B = 0.5 g_A$ and $g_{AB} = 0.1 g_A$. Each species is allowed to occupy $M = 4$ and each boson $m = 3$ SPFs both in the ML-MCTDH and in the ML-MCTDHB simulation. The dashed vertical lines indicate the first three Rabi-tunneling periods of a single particle in this double well trap.

For judging the convergence of the simulations, we plot the natural populations for different subsystems. Figure 13 shows the natural populations corresponding to the reduced density matrix of the whole species A (or B). One clearly sees that after about 25 time units three species states contribute to the total wave function with weights of the order of 89%, 10% and 1%. The fourth state contributes so little that it could be neglected without affecting the results. The natural populations of the reduced density matrix corresponding to an A and a B boson, respectively, are plotted in figure 14. Here again, we notice that the lowest natural population stays well below 1%, meaning that its natural orbital, the corresponding eigenstate of the respective reduced one-body density matrix, has only marginal influence on the result. Furthermore, we observe that two natural orbitals contribute with almost equal weight during certain time intervals. Hence, a mean field approximation would be




Figure 12: Evolution of several joint probabilities for the same setup as in figure 11: The blue (red) solid line refers to the probability of finding both A (B) bosons in the same well. The probability of detecting an A and an B boson in the same well is plotted as the green solid line. All solid lines are obtained by ML-MCTDHB, while the crosses correspond to a ML-MCTDH simulation. Again, the first three Rabi-tunneling periods are represented by the dashed vertical lines.

improper, which was to be expected for such a few-body system. Finally, we note that the ML-MCTDHB results excellently agree with the simulation performed with the well tested method ML-MCTDH.

Let us now explore a more complex tunneling scenario of three species, A, B and C, each consisting of 6 bosons. This setup both unravels interesting correlation effects and illustrates the beneficial scaling of ML-MCTDHB by introducing the extra species layer. Here, we choose $g_A(N_A - 1) = 0.2$, $g_B = 0.75 g_A$ and $g_{AB} = 0.05 \sqrt{g_A g_B}$. Furthermore, the C bosons are assumed to have no intra-species interaction, i.e. $g_C = 0$, but a weak attractive coupling to the bosons of species A and B: $g_{AC} = g_{BC} = -0.5 g_{AB}$.

From figure 15, we see that the A, B and C bosons exhibit Rabi tunneling with respect to the tunneling period. The probability amplitude for the A and the B bosons, however, decreases in the course of time, which might be a sign of equilibration. This is also supported





Figure 13: The natural populations, i.e. eigenvalues of the reduced density matrix corresponding to the whole species A (or equivalently B) are plotted versus time for the same tunneling scenario as in figure 11. The solid lines refer to the ML-MCTDHB and the crosses to the corresponding ML-MCTDH calculation.


by a low accuracy long time propagation up to 300 time units with $m = M = 3$, where we found that for $t > 100$ time units the probability amplitude for the A and the B bosons fluctuates around 0.5 with a fluctuation amplitude of less than 0.15 (data not shown). Figure 15 also shows the corresponding mean field results, which are obtained by setting $m = M = 1$. Then, the ML-MCTDHB equations of motion effectively reduce to a set of three coupled Gross-Pitaevskii equations. We clearly see that the decrease of the probability amplitude is a many-body property, not present in the mean field description. In contrast to this, the tunneling amplitude of the C bosons is not damped and its dynamics in the many-body calculation coincides with the mean field description.

A further phenomenon not captured in the mean field picture is unravelled in figure 16: The probabilities for finding two bosons of the same species in the same well oscillate between 0.5 and 1.0 with the frequency $2/T$ in the mean field calculation. In the many-body calculation, however, the probability for finding two A or B bosons in the same well features damped oscillations leading to a saturation of 0.73, which indicates a bunching tendency, while the probability of finding two C bosons stays oscillating between 0.5 and 1.0.

In order to measure the correlations within a species, we compare the joint probability



(a)



(b)

Figure 14: The dynamics of the natural populations of the reduced density matrix corresponding to an A and a B boson are shown in (a) and (b), respectively. All parameters are chosen as in figure 11. The solid lines refer to the ML-MCTDHB and the crosses to the corresponding ML-MCTDH calculation.

$P(\sigma, \sigma)$ for finding two σ bosons in the same well with the same joint probability in the case of statistical independence. If $P(\sigma; L)$ refers to the probability of finding a σ boson in the left well, the latter joint probability reads $P_{ind}(\sigma, \sigma) = [P(\sigma; L)]^2 + [1 - P(\sigma; L)]^2$. Then the ratio $f(\sigma) := P(\sigma, \sigma)/P_{ind}(\sigma, \sigma)$ measures the intra-species correlation: $f(\sigma)$ equals one if

and only if the σ bosons tunnel independently. From figure 17(a), we may infer that the ratio f stays one for the C bosons while for the A and B bosons f becomes larger than one and increases during the propagation time. Thus, the A as well as the B bosons show a bunching behaviour, which is not captured in the mean field picture, of course. Figure 17(b) shows the ratio $f(A, B) := P(A, B)/P_{ind}(A, B)$ corresponding to the joint probability of finding an A and a B boson in the same well. From this figure, we may infer that the presence of the weak attraction between the C bosons and the A (B) bosons reduces the bunching tendency of an A and a B boson being in the same well.

For discussing the convergence of the simulation, the ML-MCTDHB calculation for $m = M = 3$ is compared with the results for $m = M = 4$ in figures 15 and 16. Both the single particle and the two particle probabilities show an excellent agreement. Only for the joint probability of finding two particles of different kind in the same well, there are marginal deviations for long propagation times (not shown). Hence, we may regard the simulation as being converged. This judgement is also supported by the time evolution of the natural populations: From figure 18(a), we infer that most of the time only two natural populations significantly contribute to the reduced density matrix of the whole species A (or B or C) and, hence, to the total wave function. For times larger than 69, a third species state gains a weight over 1%. Figure 18(b) shows that the initially fully condensed state of the A bosons evolves into a two-fold fragmented state. Increasing the number of particle SPFs from $m = 3$ to 4 just leads to a reshuffling of the third-highest natural population without affecting the results. The natural populations corresponding to an B boson show a similar behaviour due to the similar intra-species interaction strengths (not shown). In contrast to this, the C bosons stay in a condensed state. A (not fully converged) long time simulation with $m = M = 3$ shows that the state of the C bosons becomes depleted by 2% due to the weak inter-species attraction (also not shown).

We finally compare the scaling of the methods MCTDH, ML-MCTDH, MCTDHB and ML-MCTDHB applied to this multi-species setup by investigating the memory consumption for storing the total wave function. This also provides us with a rough estimate for the performances of the methods. Let S be the number of species. If we denote the number of DVR grid points by n and neglect the symmetrization option as well as the possibility of primitive mode combination, $m^{SN} + SNmn$ coefficients have to be propagated in MCTDH. The ML-MCTDH expansion equivalent to our species ML-MCTDHB expansion requires




Figure 15: Tunneling dynamics of 6 A, 6 B and 6 C bosons, initially loaded into the left well, for the interaction strengths $g_A(N_A - 1) = 0.2$, $g_B = 0.75 g_A$, $g_C = 0.0$, $g_{AB} = 0.05 \sqrt{g_A g_B}$ and $g_{AC} = g_{BC} = -0.5 g_{AB}$: The probabilities for finding an A boson, a B boson and a C boson in the left well are plotted as the blue, red and green line, respectively. The solid lines refer to a ML-MCTDHB calculation with three species and three particle SPFs per species, $m = M = 3$, while the corresponding crosses are obtained for $m = M = 4$. The dotted lines show the respective mean field calculation, i.e. $m = M = 1$. The first three Rabi-tunneling periods are represented by the dashed vertical lines.

$M^S + SMm^N + SNmn$ coefficients. For a direct MCTDHB expansion, one needs $\binom{N+m-1}{m-1}^S + Smn$ coefficients. Finally, the species ML-MCTDHB ansatz consists of $M^S + SM\binom{N+m-1}{m-1} + Smn$ coefficients. Table I lists the memory consumption of the different methods for $S = 3$ species, $N = 6$ bosons per species, $n = 110$ spatial gridpoints and varied numbers of m particle and M species SPFs. If the necessary number of species SPFs is not too large with respect to the total number of bosonic configurations, $\binom{N+m-1}{m-1}$, ML-MCTDHB clearly requires less memory than all the other methods. In the case of the three species tunneling




Figure 16: For the setting of figure 15, the time evolution of the probabilities for finding two A bosons, two B bosons and two C bosons in the same well are plotted as the blue, red and green line, respectively. Solid lines correspond to a ML-MCTDHB calculation with $m = M = 3$ and crosses to $m = M = 4$, while dotted lines are the respective mean field results.

example and $m = M = 4$, the ML-MCTDHB wave function ansatz consists of two orders of magnitude less coefficients than the corresponding MCTDHB ansatz. For large numbers of species SPFs, however, a MCTDHB expansion becomes preferable with respect to the memory consumption, which can be seen from the $m = 3$ and $M = \binom{N+m-1}{m-1} = 28$ example in table I.

IV. CONCLUSION AND OUTLOOK

The methods MCTDHB and ML-MCTDH have different emphases: While MCTDHB aims at employing the bosonic particle exchange symmetry for obtaining a better performance, ML-MCTDH focuses on how to make use of system intrinsic correlations in order





Figure 17: For the setting of figure 15, the time evolution of $f(\sigma)$ (see definition in the text) is plotted for $\sigma = A, B, C$ as the blue, red and green line, respectively, in (a). Solid lines correspond to a ML-MCTDHB calculation with $m = M = 3$ and crosses to $m = M = 4$. For a mean field calculation, one finds $f(\sigma) \equiv 1.0$ (not plotted). The solid line in (b) shows the time evolution of the joint probability of finding an A and a B boson in the same well relative to that joint probability in the case of statistical independence.


The dots correspond to the same quantity in the absence of species C.

to reduce the complexity of the many-body wave function ansatz. In general however, the high flexibility of the ML-MCTDH wave function ansatz is incompatible with the generic correlations due to the bosonic exchange symmetry in a MCTDHB wave function expansion.

In this work, we have derived and applied the novel ML-MCTDHB method, which overcomes the above conceptual problems and benefits from both the flexibility of a multi-layer structure of the wave function ansatz and from an explicit consideration of the bosonic exchange symmetry. We have shown that such a symbiosis is indeed feasible for two - from a physical point of view quite natural - classes of wave function expansion schemes (and any combination thereof): (i) In species ML-MCTDHB, the total wave function of a bosonic multi-component system firstly is expanded in Hartree products made of states with each of them corresponding to a state of a whole species. Then each of these species states is expanded in permanents like in MCTDHB. (ii) In particle ML-MCTDHB, a single component bosonic system is considered with the bosons living in two or three dimensions and/or



(a)



(b)

Figure 18: The dynamics of the eigenvalues of the reduced density matrices corresponding to the whole species A (or B or C) and to a single A boson are shown in (a) and (b), respectively. The solid lines are the $m = M = 3$ ML-MCTDHB results, while crosses correspond to $m = M = 4$. The horizontal dashed line in (b) indicates an ordinate value of 0.5, i.e. the perfect twofold fragmented state. All parameters are chosen as in figure 15.

having internal degrees of freedom. There, the total wave function is expanded in terms of permanents and a ML-MCTDH expansion is applied to all the orbitals underlying those permanents. Summarizing, the bosonic exchange symmetry is exactly and efficiently taken into account in ML-MCTDHB as any state of indistinguishable bosons is expanded in permanents. The multi-layer concept is then employed for obtaining an optimized wave function

(m, M)	MCTDH	ML-MCTDH	MCTDHB	ML-MCTDHB
(1,1)	$2.0*10^3$ / 5.9	$2.0*10^3$ / 5.9	$3.3*10^2$ / 1.0	$3.3*10^2$ / 1.0
(3,1)	$3.9*10^8$ / $3.6*10^5$	$8.1*10^3$ / 7.6	$2.3*10^4$ / 21.3	$1.1*10^3$ / 1.0
(3,3)	$3.9*10^8$ / $3.1*10^5$	$1.3*10^4$ / 9.9	$2.3*10^4$ / 18.1	$1.3*10^3$ / 1.0
(3,10)	$3.9*10^8$ / $1.4*10^5$	$2.9*10^4$ / 10.2	$2.3*10^4$ / 8.1	$2.8*10^3$ / 1.0
(3,25)	$3.9*10^8$ / $2.1*10^4$	$7.6*10^5$ / 4.1	$2.3*10^4$ / 1.2	$1.9*10^4$ / 1.0
(3,27)	$3.9*10^8$ / $1.7*10^4$	$8.5*10^5$ / 3.7	$2.3*10^4$ / 1.0	$2.3*10^4$ / 1.0
(3,28)	$3.9*10^8$ / $1.5*10^4$	$8.9*10^5$ / 3.5	$2.3*10^4$ / 0.9	$2.5*10^4$ / 1.0
(4,1)	$6.9*10^{11}$ / $4.4*10^7$	$2.0*10^4$ / 12.8	$5.9*10^5$ / 377.6	$1.6*10^3$ / 1.0
(4,3)	$6.9*10^{11}$ / $3.3*10^7$	$4.4*10^4$ / 21.3	$5.9*10^5$ / 282.5	$2.1*10^3$ / 1.0
(4,4)	$6.9*10^{11}$ / $2.9*10^7$	$5.7*10^4$ / 23.9	$5.9*10^5$ / 248.3	$2.4*10^3$ / 1.0

Table I: The scaling of the methods MCTDH, ML-MCTDH, MCTDHB and ML-MCTDHB is compared for the case of three species, each consisting of 6 bosons. The number of particle SPFs, m , and (for the multi-layer methods) the number of species SPFs, M , are varied. The spatial grid is assumed to consist of $n = 110$ points. Each table entry contains the number of coefficients needed for the wave function expansion of the respective method and its ratio with respect to the number of ML-MCTDHB coefficients.

ansatz guided by the correlations between different species (intra- versus inter-species correlations) or between different spatial directions (e.g. in quasi two- or quasi three-dimensional systems) or internal degrees of freedom. By means of a mixture tunneling scenario, we have illustrated the beneficial scaling of ML-MCTDHB. Like in all MCTDH type methods, the convergence of a simulation with respect to the number of provided states in the wave function ansatz can be judged by the eigenvalue distributions of the reduced density matrices corresponding to different subsystems.

The equations of motion of ML-MCTDHB can be derived by employing the Dirac-Frenkel variational principle ensuring that at any instant in time the propagated wave function remains variationally optimal within the class of wave functions given by the ansatz. Here, we presented an alternative but equivalent derivation, which uses the ML-MCTDH equations of motion as a starting point. Moreover, we have proven that all MCTDH type methods preserve an arbitrary symmetry of a given system in the sense explicated in section II C 2

as long as the corresponding symmetry operation can be written as a product of symmetry operations acting on only a single physical DOF. A generalization to genuine “many-body” symmetry operations acting on more than one physical DOF remains an open issue.

We have implemented ML-MCTDHB on the basis of the ML-MCTDH code [16] in such a general way that, in principle, we can deal with an arbitrary number of species in arbitrary dimensions - only limited by the number of states needed for a converged simulation, which depends on the system details, of course. Furthermore, the code can in principle also deal with hybrid systems such as a single ion coupled to an environment of indistinguishable bosons as long as all the interactions can be efficiently brought into the POTFIT product form.

Furthermore, we have shown the first applications of the ML-MCTDHB method in terms of various correlated tunneling scenarios of a double well potential. Particularly for a setup with three species named A, B, C, where all interactions have been chosen to be repulsive except for the C bosons, which have no intra-species interaction but weakly attract bosons of the other two species, we have observed the emergence of intra- and inter-species correlations and signatures of equilibration: The probability of finding an boson of kind A (B) in the left well tends to equilibrate at a value of about 0.5, which is not captured in a mean field description at all. Similarly, the probability of finding two A (B) bosons in the same well tends to equilibrate at a value of 0.7. Such correlations can intrinsically not be described by the mean field ansatz. A more careful investigation has shown that the interactions lead to a bunching tendency of the A (B) bosons, which is increasing in time. We have shown that the weak attraction between the C and A (B) bosons reduces the slight bunching tendency of finding an A and a B boson in the same well. To summarize, there are various interaction induced correlation effects in such a three species tunneling setup, which can be studied systematically in further works. In particular, a discrimination of classical from quantum correlation appears to be a challenging perspective.

Finally, we note that both conceptually and practically it is relatively straightforward to generalize the presented method to fermionic systems, i.e. to ML-MCTDHF, or to mixed bosonic-fermionic systems, i.e. to ML-MCTDHFHB: According to the presented correspondence principle, one could start with an appropriate ML-MCTDH expansion in which all indistinguishable particles of one kind are grouped into a species node. The ML-MCTDH expansion of each of these species nodes would then have to be reformulated in the sec-

and quantization picture, i.e. in terms of permanents or Slater determinants depending on whether the species is bosonic or fermionic. This approach is in particular compatible with the present ML-MCTDH code [16] and our ML-MCTDHB extension.

ACKNOWLEDGMENTS

The authors would like to thank Hans-Dieter Meyer and Jan Stockhofe for fruitful discussions on MCTDH methods and symmetry conservation. Particularly, the authors would like to thank Jan Stockhofe for the DVR implementation of the ML-MCTDHB code. S.K. gratefully acknowledges financial support by the German National Academic Foundation. L.C. and P.S. gratefully acknowledge funding by the Deutsche Forschungsgemeinschaft in the framework of the SFB 925 “Light induced dynamics and control of correlated quantum systems”.

Appendix A: Proof of Symmetry Conservation in MCTDH and its Derivatives

Here, we sketch the proof that MCTDH (and all its extensions) preserves both the symmetry of the initial many-body wave function and the initial SPFs as explicated in section II C 2. According to a standard argument, the Hamiltonian matrix $\langle I|\hat{H}|J\rangle$ in (11) does not allow transitions from Hartree products $|J\rangle$ with $\hat{G}|J\rangle = e^{i\Theta'}|J\rangle$ to Hartree products $|I\rangle$ with $\hat{G}|I\rangle = e^{i\Theta''}|I\rangle$ and $\Theta' \bmod 2\pi \neq \Theta'' \bmod 2\pi$, when propagating the coefficients for a small time step Δt , i.e. $\langle I|\hat{H}|J\rangle$ vanishes for such Hartree products. Hence, we are left to show that the SPFs preserve their symmetry up to $O(\Delta t^2)$ when propagating also them for a small time step. We note that the mean field operator matrices of a Hamiltonian containing at most F -body terms can be expressed by the F -body reduced density matrices and one-body operators. If we group the initial SPFs $|\phi_j(0)\rangle$ into classes of same $\theta_j \bmod 2\pi$ (cf. section II C 2), the F -body reduced density matrices and the inverses of the one-body reduced density matrices in (12) are initially block diagonal. With this, one can show $i\hat{g}_j\partial_t|\phi_j(t)\rangle = ie^{i\theta_j}\partial_t|\phi_j(t)\rangle$ at $t = 0$. The same arguments hold for any later time steps as long as Δt is small enough and the symmetry conservation becomes exact for $\Delta t \rightarrow 0$.

Appendix B: Equations of motion for species-ML MCTDHB

In this appendix, we demonstrate how to construct the equations of motion for the coefficients of each layer in species multi-layer MCTDHB. Firstly, let us state the recursive formulas for calculating the reduced density matrices in equations (36). The density matrices of the species layer are computed in the well known MCTDH manner:

$$\rho_{nm}^{2;\kappa} = \sum_{J^\kappa} \left(A_{J_n^\kappa}^{2;\kappa} \right)^* A_{J_m^\kappa}^{2;\kappa}, \quad (\text{B1})$$

where J^κ indicates a summation over all species SPFs indices j_σ but j_κ . J_n^κ refers to the multi-index $(j_1, \dots, j_{\kappa-1}, n, j_{\kappa+1}, \dots, j_S)$. For calculating the reduced density matrix on the physical layer, we have to evaluate the SHFs of a bosonic particle node at first:

$$|\Psi_i^{3;\kappa}\rangle := \frac{\hat{a}_{\kappa,i}}{\sqrt{N_\kappa}} |\Psi\rangle = \sum_j |\Psi_j^{2;\kappa}\rangle \left(\sum_{\vec{n}|N_\kappa-1} A_{j;\vec{n}+\hat{i}}^{2;\kappa} \sqrt{(n_i+1)/N_\kappa} |\vec{n}\rangle_\kappa \right), \quad (\text{B2})$$

which leads to:

$$\rho_{ij}^{3;\kappa} = \frac{1}{N_\kappa} \sum_{k,l=1}^{M_\kappa} \rho_{kl}^{2;\kappa} \sum_{\vec{n}|N_\kappa-1} Q_{\vec{n}}(i, j) \left(A_{k;\vec{n}+\hat{i}}^{2;\kappa} \right)^* A_{l;\vec{n}+\hat{j}}^{2;\kappa}, \quad (\text{B3})$$

with $Q_{\vec{n}}(i, j) = \sqrt{(n_i+1)(n_j+1)}$.

In order to derive formulas for the mean field operators, we should also express the Hamiltonian (33) in the second quantization language:

$$\begin{aligned} \hat{H}_\kappa &= \sum_{i,j=1}^{m_\kappa} \langle \phi_i^{3;\kappa} | \hat{h}^\kappa | \phi_j^{3;\kappa} \rangle \hat{a}_{\kappa,i}^\dagger \hat{a}_{\kappa,j}, \\ \hat{V}_\kappa &= \frac{1}{2} \sum_\nu \sum_{p,q,i,j=1}^{m_\kappa} C_\nu^\kappa \langle \phi_p^{3;\kappa} | \hat{v}_1^{\kappa,\nu} | \phi_i^{3;\kappa} \rangle \langle \phi_q^{3;\kappa} | \hat{v}_2^{\kappa,\nu} | \phi_j^{3;\kappa} \rangle \hat{a}_{\kappa,p}^\dagger \hat{a}_{\kappa,q}^\dagger \hat{a}_{\kappa,i} \hat{a}_{\kappa,j}, \\ \hat{W}_{\kappa,\sigma} &= \sum_\nu \sum_{p,q=1}^{m_\kappa} \sum_{i,j=1}^{m_\sigma} D_\nu^{\kappa,\sigma} \langle \phi_p^{3;\kappa} | \hat{w}^{\kappa,\nu} | \phi_q^{3;\kappa} \rangle \langle \phi_i^{3;\sigma} | \hat{w}^{\sigma,\nu} | \phi_j^{3;\sigma} \rangle \hat{a}_{\kappa,p}^\dagger \hat{a}_{\kappa,q} \hat{a}_{\sigma,i}^\dagger \hat{a}_{\sigma,j}. \end{aligned} \quad (\text{B4})$$

Here, we have introduced the one-body operators $\hat{h}^\kappa \equiv \hat{h}_1^\kappa$ and $\hat{w}^{\eta,\nu} \equiv \hat{w}_1^{\eta,\nu}$ to simplify the notation. Please note that the instantaneous matrix elements occurring in (B4) have to be evaluated by means of the PBFs according to (32).

Going from the physical layer upwards, we construct the mean field operator matrices and state the equations of motion layer by layer. Due to the presence of the projector $\hat{P}^{\alpha;\kappa}$ and the projection on $|C\rangle$ in (36), we have to calculate the matrix elements $\langle F | \langle \Psi_n^{\alpha;\kappa} | \hat{H} | \Psi_m^{\alpha;\kappa} \rangle | E \rangle$,

where $|E\rangle$ and $|F\rangle$ are two PBFs or permanents for the physical ($\alpha = 3$) and the species ($\alpha = 2$) layer, respectively. We are allowed to treat the one-body, intra-species and inter-species interaction terms separately. Furthermore, we may employ the factorization of the mean field operator matrix into an operator and a scalar prefactor, the mean field matrix, (14) due to the product structure (34).

First, we investigate the physical layer. But before diving into the concrete calculations, a word of caution is in order here: In (B2), we have introduced the SHF of a bosonic physical DOF x_κ , which clearly contains $N_\kappa - 1$ bosons of species κ . However, the mean field operator $\langle \Psi_n^{3;\kappa} | \hat{H} | \Psi_m^{3;\kappa} \rangle$ is in fact an operator acting on single particle states. So one must not plug in the Hamiltonian in second quantization representation (B4) directly, rather one should use the first quantization representation and reformulate the result in terms the second quantization language according to the correspondence principle (cf. also footnote 9).

The peculiarity of the one-body terms $\hat{H}_\kappa$ is that they turn out to be uncorrelated in the sense that $\langle \Psi_n^{3;\kappa} | \hat{H}_\kappa | \Psi_m^{3;\kappa} \rangle$ equals $\hat{h}_\kappa \rho_{nm}^{3;\kappa}$ plus scalar terms which are killed by the projector. Hence, the scalar factor of the surviving term will cancel the inverse density matrix in (36). In contrast to this, the mean field operators of the interaction terms show non-trivial mean field matrices:

$$\begin{aligned} \langle \Psi_n^{3;\kappa} | \hat{V}_\eta | \Psi_m^{3;\kappa} \rangle &= \delta_{\eta\kappa} (N_\kappa - 1) \sum_\nu C_\nu^\kappa \langle \Psi_n^{3;\kappa} | \hat{v}_2^{\kappa,\nu} | \Psi_m^{3;\kappa} \rangle \hat{v}_1^{\kappa,\nu} + \text{scalar terms}, \\ \langle \Psi_n^{3;\kappa} | \hat{W}_{\eta,\sigma} | \Psi_m^{3;\kappa} \rangle &= \sum_\nu D_\nu^{\eta,\sigma} (\delta_{\eta\kappa} N_\sigma \langle \Psi_n^{3;\kappa} | \hat{w}^{\sigma,\nu} | \Psi_m^{3;\kappa} \rangle + \\ &\quad + \delta_{\sigma\kappa} N_\eta \langle \Psi_n^{3;\kappa} | \hat{w}^{\eta,\nu} | \Psi_m^{3;\kappa} \rangle) \hat{w}^{\kappa,\nu} + \text{scalar terms}. \end{aligned} \quad (\text{B5})$$

We again do not have to care about the scalar terms since they will be killed by the projector. Probably the easiest way for arriving at the above expressions is to think in the first quantization picture and to interpret the SPF $|\phi_i^{3;\kappa}\rangle$ of the bosonic physical DOF x_κ as a SPF of the “first” κ boson in an - according to the correspondence principle - equivalent ML-MCTDH expansion of the total wave function. Then the SHFs $|\Psi_k^{3;\kappa}\rangle$ describe states corresponding to the “second” up to the “ N_κ -th” boson of the species κ and all the bosons of the other species.

Now we may interpret the operators in the mean field matrices as one-body operators in

second quantization language, which enables us to apply the relation (B2) directly:

$$\begin{aligned}
\langle v_2^{\kappa,\nu} \rangle_{nm}^{3;\kappa} &:= (N_\kappa - 1) \langle \Psi_n^{3;\kappa} | \hat{v}_2^{\kappa,\nu} | \Psi_m^{3;\kappa} \rangle \\
&= \sum_{i,j=1}^{m_\kappa} \langle \phi_i^{3;\kappa} | \hat{v}_2^{\kappa,\nu} | \phi_j^{3;\kappa} \rangle \langle \Psi_n^{3;\kappa} | \hat{a}_{\kappa,i}^\dagger \hat{a}_{\kappa,j} | \Psi_m^{3;\kappa} \rangle, \\
\langle w^{\sigma,\nu} \rangle_{nm}^{3;\kappa} &:= N_\sigma \langle \Psi_n^{3;\kappa} | \hat{w}^{\sigma,\nu} | \Psi_m^{3;\kappa} \rangle = \sum_{i,j=1}^{m_\sigma} \langle \phi_i^{3;\sigma} | \hat{w}^{\sigma,\nu} | \phi_j^{3;\sigma} \rangle \langle \Psi_n^{3;\kappa} | \hat{a}_{\sigma,i}^\dagger \hat{a}_{\sigma,j} | \Psi_m^{3;\kappa} \rangle.
\end{aligned} \tag{B6}$$

So we are left to evaluate $\langle \Psi_n^{3;\kappa} | \hat{a}_{\kappa,i}^\dagger \hat{a}_{\kappa,j} | \Psi_m^{3;\kappa} \rangle$ and $\langle \Psi_n^{3;\kappa} | \hat{a}_{\sigma,i}^\dagger \hat{a}_{\sigma,j} | \Psi_m^{3;\kappa} \rangle$, which are essentially the reduced two-body intra-species density matrix of species κ and the reduced two-body inter-species density matrix of species κ and σ , respectively. Thus, those quantities can also be employed for a physical interpretation of a ML-MCTDHB simulation.

$$\begin{aligned}
\langle \Psi_n^{3;\kappa} | \hat{a}_{\kappa,i}^\dagger \hat{a}_{\kappa,j} | \Psi_m^{3;\kappa} \rangle &= \frac{1}{N_\kappa} \sum_{r,s} \rho_{rs}^{2;\kappa} \sum_{\vec{l}|N_\kappa-2} \left(A_{r;\vec{l}+\hat{n}+\hat{i}}^{2;\kappa} \right)^* P_{\vec{l}}(n, i) \\
&\quad \times A_{s;\vec{l}+\hat{m}+\hat{j}}^{2;\kappa} P_{\vec{l}}(m, j), \\
\langle \Psi_n^{3;\kappa} | \hat{a}_{\sigma,i}^\dagger \hat{a}_{\sigma,j} | \Psi_m^{3;\kappa} \rangle &= \frac{1}{N_\kappa} \sum_{r,s,u,v} \eta_{ru;sv}^{2;\kappa\sigma} \sum_{\vec{l}|N_\kappa-1} \left(A_{r;\vec{l}+\hat{n}}^{2;\kappa} \right)^* A_{s;\vec{l}+\hat{m}}^{2;\kappa} Q_{\vec{l}}(n, m) \\
&\quad \times \sum_{\vec{o}|N_\sigma-1} \left(A_{u;\vec{o}+\hat{i}}^{2;\sigma} \right)^* A_{v;\vec{o}+\hat{j}}^{2;\sigma} Q_{\vec{o}}(u, v),
\end{aligned} \tag{B7}$$

where $P_{\vec{l}}(n, i) = \sqrt{(l_n + \delta_{ni} + 1)(l_i + 1)}$ and $\eta_{ru;sv}^{2;\kappa\sigma}$ denoting the reduced two-body inter-species density matrix in the respective species SPF representation:

$$\eta_{ru;sv}^{2;\kappa\sigma} = \sum_{J^{\kappa\sigma}} \left(A_{ru}^{1;\kappa\sigma} \right)^* A_{sv}^{1;\kappa\sigma}. \tag{B8}$$

Here, $J^{\kappa\sigma}$ refers to a summation over all indices but those of κ and σ and $J_{ru}^{\kappa\sigma}$ denotes the multi-index $J^{\kappa\sigma}$ with κ being in the r -th and σ being u -th species SPF. Hence, we arrive at the final result for the equations of motion of the physical layer:

$$\begin{aligned}
i\partial_t A_{i;r}^{3;\kappa} &= \sum_s \langle x_r^\kappa | (1 - \hat{P}^{3;\kappa}) \hat{h}_\kappa | x_s^\kappa \rangle A_{i;s}^{3;\kappa} + \\
&\quad + \sum_s \sum_\nu C_\nu^\kappa \langle x_r^\kappa | (1 - \hat{P}^{3;\kappa}) \hat{v}_1^{\kappa,\nu} | x_s^\kappa \rangle \sum_{n,m} (\rho^{3;\kappa})_{im}^{-1} \langle v_2^{\kappa,\nu} \rangle_{mn}^{3;\kappa} A_{n;s}^{3;\kappa} \\
&\quad + \sum_s \sum_{\sigma \neq \kappa} \sum_\nu D_\nu^{\kappa,\sigma} \langle x_r^\kappa | (1 - \hat{P}^{3;\kappa}) \hat{w}^{\kappa,\nu} | x_s^\kappa \rangle \sum_{n,m} (\rho^{3;\kappa})_{im}^{-1} \langle w^{\sigma,\nu} \rangle_{mn}^{3;\kappa} A_{n;s}^{3;\kappa}.
\end{aligned} \tag{B9}$$

Please note that the projector $\hat{P}^{3;\kappa} = \sum_i |\phi_i^{3;\kappa}\rangle \langle \phi_i^{3;\kappa}|$ also has to be expressed in terms of the PBFs by means of (32). Furthermore, we have assumed that $D_\nu^{\sigma,\kappa} = D_\nu^{\kappa,\sigma}$ without loss of generality.

Next we turn to the ingredients of the equations of motion for the species layer. In this layer, the SHFs $|\Psi_i^{2;\kappa}\rangle$ describe states of all species but κ . Following the same logic as above, the one-body and intra-species interaction terms, $\hat{H}_\kappa$ and $\hat{V}_\kappa$, respectively, are uncorrelated terms on the species layer. Thus, we are left to determine the mean field operators corresponding to $\hat{W}_{\eta,\sigma}$, which are only non-trivial for either η or σ equal to κ . Here, we evaluate the mean field operators for $\eta = \kappa$ after separating them into mean field matrices and operator valued terms (cf. (14)):

$$\langle \Psi_n^{2;\kappa} | \hat{W}_{\kappa,\sigma} | \Psi_m^{2;\kappa} \rangle = \sum_\nu D_\nu^{\kappa,\sigma} \langle W_\nu^{\kappa,\sigma} \rangle_{nm}^{3;\kappa} \hat{w}_\nu^{2;\kappa}. \quad (\text{B10})$$

By employing the second quantization representation (B4), we can state the operator valued factor:

$$\hat{w}_\nu^{2;\kappa} = \sum_{p,q=1}^{m_\kappa} \langle \phi_p^{3;\kappa} | \hat{w}^{\kappa,\nu} | \phi_q^{3;\kappa} \rangle \hat{a}_{\kappa,p}^\dagger \hat{a}_{\kappa,q} \quad (\text{B11})$$

as well as the mean field matrices:

$$\langle W_\nu^{\kappa,\sigma} \rangle_{nm}^{3;\kappa} = \sum_{i,j=1}^{m_\sigma} \langle \phi_i^{3;\sigma} | \hat{w}^{\sigma,\nu} | \phi_j^{3;\sigma} \rangle \langle \Psi_n^{2;\kappa} | \hat{a}_{\sigma,i}^\dagger \hat{a}_{\sigma,j} | \Psi_m^{2;\kappa} \rangle. \quad (\text{B12})$$

Inserting the definition of the SHF leads to the following:

$$\langle \Psi_n^{2;\kappa} | \hat{a}_{\sigma,i}^\dagger \hat{a}_{\sigma,j} | \Psi_m^{2;\kappa} \rangle = \sum_{r,s} \eta_{nr;ms}^{2;\kappa\sigma} \sum_{\vec{l}|N_\sigma-1} Q_{\vec{l}}(i,j) \left(A_{r;\vec{l}+i}^{2;\sigma} \right)^* A_{s;\vec{l}+j}^{2;\sigma}. \quad (\text{B13})$$

So the equations of motion for the species layer read:

$$\begin{aligned} i\partial_t A_{i;\vec{n}}^{2;\kappa} &= \sum_{\vec{m}|N_\kappa} \langle \vec{n} | (1 - \hat{P}^{2;\kappa}) (\hat{H}_\kappa + \hat{V}_\kappa) | \vec{m} \rangle A_{i;\vec{m}}^{2;\kappa} + \\ &+ \sum_{\sigma \neq \kappa} \sum_\nu D_\nu^{\kappa,\sigma} \sum_{\vec{m}|N_\kappa} \langle \vec{n} | (1 - \hat{P}^{2;\kappa}) \hat{w}_\nu^{2;\kappa} | \vec{m} \rangle \sum_{r,s} (\rho^{2;\kappa})_{ir}^{-1} \langle W_\nu^{\kappa,\sigma} \rangle_{rs}^{3;\kappa} A_{s;\vec{m}}^{2;\kappa}. \end{aligned} \quad (\text{B14})$$

Again, the projector $\hat{P}^{2;\kappa}$ has to be explicated by means of (31). Besides, one has to evaluate the action of the operators $\hat{H}_\kappa$, $\hat{V}_\kappa$ and $\hat{w}_\nu^{2;\kappa}$ on the species SPF $|\phi_n^{2;\kappa}\rangle$. This is straightforward to do when employing the second quantization representations (B4), (B11) and, here, we only state how the archetypes of a one- and of a two-body¹⁰ operator act on such a species

¹⁰ Here, we refer to only intra-species interaction terms as two-body operators in contrast to inter-species interaction terms, which we regard as products of intra-species one-body operators (cf. (B4)).

SPF:

$$\begin{aligned}\hat{a}_{\kappa,i}^\dagger \hat{a}_{\kappa,j} |\phi_n^{2;\kappa}\rangle &= \sum_{\vec{l}|N_\kappa-1} Q_{\vec{l}}(i,j) A_{n;\vec{l}+\hat{j}}^{2;\kappa} |\vec{l} + \hat{i}\rangle, \\ \hat{a}_{\kappa,p}^\dagger \hat{a}_{\kappa,q}^\dagger \hat{a}_{\kappa,i} \hat{a}_{\kappa,j} |\phi_n^{2;\kappa}\rangle &= \sum_{\vec{l}|N_\kappa-2} P_{\vec{l}}(p,q) P_{\vec{l}}(i,j) A_{n;\vec{l}+\hat{i}+\hat{j}}^{2;\kappa} |\vec{l} + \hat{p} + \hat{q}\rangle.\end{aligned}\tag{B15}$$

When plugging (B4) and (B15) into (B14), we arrive at the final equations of motion. Finally, we turn to the top layer equation of motion:

$$\begin{aligned}i\partial_t A_I^1 &= \sum_{\kappa} \sum_{p=1}^{M_\kappa} \langle \phi_{i_\kappa}^{2;\kappa} | (\hat{H}_\kappa + \hat{V}_\kappa) | \phi_p^{2;\kappa} \rangle A_{I_p^\kappa}^1 + \\ &+ \sum_{\kappa < \sigma} \sum_{p=1}^{M_\kappa} \sum_{q=1}^{M_\sigma} \langle \phi_{i_\kappa}^{2;\kappa} | \langle \phi_{i_\sigma}^{2;\sigma} | \hat{W}_{\kappa,\sigma} | \phi_p^{2;\kappa} \rangle | \phi_q^{2;\sigma} \rangle A_{I_{p,q}^{\kappa,\sigma}}^1.\end{aligned}\tag{B16}$$

The occurring mean field matrices can be calculated with the help of (B15) and the Hamiltonian in second quantization language (B4). Here, we only state the result for the mean field matrix corresponding to the intra-species interaction:

$$\begin{aligned}\langle \phi_n^{2;\kappa} | \hat{V}_\kappa | \phi_m^{2;\kappa} \rangle &= \frac{1}{2} \sum_{p,q,i,j=1}^{m_\kappa} \sum_{\nu} C_\nu^\kappa \langle \phi_p^{3;\kappa} | \hat{v}_1^{\kappa,\nu} | \phi_i^{3;\kappa} \rangle \langle \phi_q^{3;\kappa} | \hat{v}_2^{\kappa,\nu} | \phi_j^{3;\kappa} \rangle \\ &\sum_{\vec{l}|N_\kappa-2} P_{\vec{l}}(p,q) P_{\vec{l}}(i,j) \left(A_{n;\vec{l}+\hat{p}+\hat{q}}^{2;\kappa} \right)^* A_{m;\vec{l}+\hat{i}+\hat{j}}^{2;\kappa}.\end{aligned}\tag{B17}$$

Appendix C: Comments on the Implementation

In several of the equations above, there are summations over bosonic number state configurations. In a computer program, these sums can be performed conveniently in the mathematical framework of combinatorial number systems [57] as pointed out in [25]. The idea is to map a number state $\vec{n}$ to a single integer, which serves as a unique address for picking the corresponding coefficient, say $A_{i;\vec{n}}^{2;\kappa}$, out of an array. In [25], a one- or two-body operator is applied to a bosonic state as follows: A nested loop over all occupation numbers $n_1, \dots, n_m$ is performed with the constraint that $\sum_i n_i$ equals the number of bosons (of one kind). Applying a one- or two-body operator results in removing and adding bosons in certain orbitals. The “scattered” number state can then be mapped to a single integer and, thus, the action of a one- or two-body operator reduces to re-addressing the number states (and multiplying the respective coefficient with a known prefactor).

In our ML-MCTDHB code, we employ a scheme which is different in two aspects: First of all, we would like to note that we have reduced the summation over all N particle number states to a summation over all $N - 1$ ($N - 2$) particle configurations for all one- (two-) body operator related quantities. This reduction improves the performance due to the binomial scaling of the number of configurations with the number of bosons. Secondly, our code calculates several index mapping tables before the actual simulation: (i) There is a table from which, for every integer address I of a number state, the corresponding occupation number $n_1, \dots, n_m$ can be read off. Hence we can avoid a nested looping over the occupation numbers n_i by simply looping over the integer addresses I . (ii) For any integer address I corresponding to a $N - 1$ particle number state $\vec{l}$, we store the integer address I' corresponding to the N particle number state $\vec{l} + \hat{i}$ for any $i = 1, \dots, m$ in another table. (iii) Similarly, for any integer address I corresponding to an $N - 2$ particle number state $\vec{l}$, we store the integer address I' corresponding to the N particle number state $\vec{l} + \hat{i} + \hat{j}$ with $i, j = 1, \dots, m$. As the formulas like (B15) indicate, applying a one- or two-body operator to a bosonic state effectively means finding the correct integer addresses of the number states from which and to which the annihilation and creation operators scatter. The tables (ii) and (iii) allow for a extremely fast and efficient identification of these addresses, while table (i) is used for reading off the underlying occupation numbers needed for the prefactors.

Finally, we would like to comment about interaction potentials which are not given in the POTFIT form. Any potential can be brought into POTFIT form [39, 40]. In general, such a representation requires as many POTFIT terms as there are gridpoints but often only a few terms are really needed for a good approximation of the potential, which improves the performance. If, however, many POTFIT terms are needed, this can slow down the simulation and it can be beneficial to work with evolution equations which are not based on a potential of the POTFIT form. For such a potential, the equations of motion can be derived by starting with (B9, B14, B16) and then performing the summation over the POTFIT terms first and foremost, which recovers the interaction potential in its “original” form. As a consequence, one cannot split the mean field operators into scalar valued mean fields and operator valued factors anymore, which breaks the recursive structure of a multi-layer expansion and leads to more complicated objects. In particular, this procedure is straightforward for the contact potential in one dimension: After inserting the exact POTFIT representation $\delta(x_1 - x_2) = \int dy \delta(x_1 - y) \delta(x_2 - y)$ into the equations of motion, the summation

over the POTFIT terms, i.e. the integral over y , can be easily performed and one arrives at the ML-MCTDHB equations of motion without the product form assumption. We have observed a speed up by one order of magnitude and sometimes even more for this efficient implementation of the contact interaction in comparison to the equations of motion with a POTFIT representation containing as many terms as there are PBFs.

-
- [1] K.B. Davis, M.O. Mewes, M.R. Andrews, N.J.v. Druten, D.S. Durfee, D.M. Kurn and W. Ketterle, *Phys. Rev. Lett.* **75**, 3969 (1995).
 - [2] M.H. Anderson, J.R. Ensher, M.R. Matthews, C.E. Wieman and E.A. Cornell, *Science* **269**, 198 (1995).
 - [3] L. Pitaevskii and S. Stringari, *Bose-Einstein Condensation*, Oxford Science Publication, Oxford Univ. Press, (Oxford, 2003).
 - [4] I. Bloch, J. Dalibard and W. Zwerger, *Rev. Mod. Phys.* **80**, 885 (2008).
 - [5] P. Soltan-Panahi, J. Struck, P. Hauke, A. Bick, W. Plenkers, G. Meineke, C. Becker, P. Windpassinger, M. Lewenstein and K. Sengstock, *Nat. Phys.* **7**, 424 (2011).
 - [6] B. Paredes, A. Widera, V. Murg, O. Mandel, S. Fölling, I. Cirac, G.V. Shlyapnikov, T.W. Hänsch and I. Bloch, *Nature* **429**, 277 (2004).
 - [7] G. Zürn, F. Serwane, T. Lompe, A.N. Wenz, M.G. Ries, J.E. Bohn and S. Jochim, *Phys. Rev. Lett.* **108**, 075303 (2012).
 - [8] M.H. Beck, A. Jäckle, G.A. Worth and H.-D. Meyer, *Phys. Rep.* **324**, 1 (2000).
 - [9] H.-D. Meyer, U. Manthe and L.S. Cederbaum, *Chem. Phys. Lett.* **165**, 73 (1990).
 - [10] H.-D. Meyer, *WIREs Comp. Mol. Sci.* **2**, 351 (2012).
 - [11] *Multidimensional Quantum Dynamics: MCTDH Theory and Applications*, ed. by H.-D. Meyer, F. Gatti and G.A. Worth, WILEY-VCH (2009).
 - [12] S. Zöllner, H.-D. Meyer, and P. Schmelcher, *Phys. Rev. Lett.* **100**, 040401 (2008).
 - [13] L. Cao, I. Brouzos, S. Zöllner and P. Schmelcher, *New J. Phys.* **13**, 033032 (2011).
 - [14] H. Wang and M. Thoss, *J. Chem. Phys.* **119**, 1289 (2003).
 - [15] U. Manthe, *J. Chem. Phys.* **128**, 164116 (2008).
 - [16] O. Vendrell and H.-D. Meyer, *J. Chem. Phys.* **134**, 044135 (2011).
 - [17] H. Wang and M. Thoss, *J. Chem. Phys.* **124**, 034114 (2006).

- [18] I. Kondov, M. Cížek, C. Benesch, H. Wang, and M. Thoss, *J. Phys. Chem. C* **111**, 11970 (2007).
- [19] H. Wang and M. Thoss, *New J. Phys.* **10**, 115005 (2008).
- [20] H. Wang and J. Shao, *J. Chem. Phys.* **137**, 22A504 (2012).
- [21] T. Hammer and U. Manthe, *J. Chem. Phys.* **134**, 224305 (2011).
- [22] T. Hammer and U. Manthe, *J. Chem. Phys.* **136**, 054105 (2012).
- [23] A.I. Streltsov, O.E. Alon and L.S. Cederbaum, *Phys. Rev. Lett.* **99**, 030402 (2007).
- [24] O.E. Alon, A.I. Streltsov and L.S. Cederbaum, *Phys. Rev. A* **77**, 033613 (2008).
- [25] A.I. Streltsov, O.E. Alon and L.S. Cederbaum, *Phys. Rev. A* **81**, 022124 (2010).
- [26] J. Zanghellini, M. Kitzler, C. Fabian, T. Brabec and A. Scrinzi, *Laser Phys.* **13**, 1064 (2003).
- [27] G. Jordan, J. Caillat, C. Ede and A. Scrinzi, *J. Phys. B* **39**, S341 (2006).
- [28] M. Nest, R. Padmanaban and P. Saalfrank, *J. Chem. Phys.* **126**, 214106 (2007).
- [29] D. Hochstuhl and M. Bonitz, *J. Chem. Phys.* **134**, 084106 (2011).
- [30] D.J. Haxton, K.V. Lawler and C.W. McCurdy, *Phys. Rev. A* **86**, 013406 (2012).
- [31] K. Sakmann, A.I. Streltsov, O.E. Alon and L.S. Cederbaum, *Phys. Rev. Lett.* **103**, 220601 (2009).
- [32] A.I. Streltsov, O.E. Alon and L.S. Cederbaum, *Phys. Rev. Lett.* **106**, 240401 (2011).
- [33] O.E. Alon, A.I. Streltsov, K. Sakmann, A.U.J. Lode, J. Grond and L.S. Cederbaum, *Chem. Phys.* **401**, 2 (2012).
- [34] O.E. Alon, A.I. Streltsov and L.S. Cederbaum, *Phys. Rev. A* **79**, 022503 (2009).
- [35] H. Wang and M. Thoss, *J. Chem. Phys.* **131**, 024114 (2009).
- [36] H. Wang, I. Pshenichnyuk, R. Härtle and M. Thoss, *J. Chem. Phys.* **135**, 244506 (2011).
- [37] K.F. Albrecht, H. Wang, L. Mühlbacher, M. Thoss and A. Komnik, *Phys. Rev. B* **86**, 081412 (2012).
- [38] H.-D. Meyer and G.A. Worth, *Theor. Chem. Acc.* **109**, 251 (2003).
- [39] A. Jäckle and H.-D. Meyer, *J. Chem. Phys.* **104**, 7974 (1996).
- [40] A. Jäckle and H.-D. Meyer, *J. Chem. Phys.* **109**, 3772 (1998).
- [41] P.A.M. Dirac, *Proc. Cambridge Philos. Soc.* **26**, 376 (1930).
- [42] J. Frenkel, *Wave Mechanics*, Clarendon Press, Oxford, 1934.
- [43] A.D. McLachlan, *Mol. Phys.* **8**, 39 (1964).
- [44] J. Broeckhove, L. Lathouwers, E. Kesteloot and P. Van Leuven, *Chem. Phys. Lett.* **149**, 547

- (1988).
- [45] J.C. Light and T. Carrington, *Adv. Chem. Phys.* **114**, 263 (2000).
 - [46] A.C. Hindmarsh, *ODEPACK, a Systematized Collection of ODE Solvers, Scientific Computing*, ed. by R.S. Stepleman et al., North-Holland, Amsterdam, pp. 55-64 (1983).
 - [47] E. Hairer, S.P. Nørsett and G. Wanner, *Solving Ordinary Differential Equations I. Nonstiff Problems*, 2nd edition, Springer Series in Computational Mathematics, Springer-Verlag (1993).
 - [48] G.A. Worth, M.H. Beck, A. Jäckle and H.-D. Meyer H-D, The MCTDH Package, Version 8.2, (2000). H-D. Meyer, Version 8.3 (2002), Version 8.4 (2007). See <http://mctdh.uni-hd.de/>.
 - [49] A. Smerzi, S. Fantoni, S. Giovanazzi and S.R. Shenoy, *Phys. Rev. Lett.* **79**, 4950 (1997).
 - [50] G. J. Milburn, J. Corney, E. M. Wright and D. F. Walls, *Phys. Rev. A* **55**, 4318 (1997).
 - [51] E. Boukobza, M. Chuchem, D. Cohen and A. Vardi, *Phys. Rev. Lett.* **102**, 180403 (2009).
 - [52] M.T. Martinez, A. Posazhennikova, and J. Kroha, *Phys. Rev. Lett.* **103**, 105302 (2009).
 - [53] M. Albiez, R. Gati, J. Fölling, S. Hunsmann, M. Cristiani and M.K. Oberthaler, *Phys. Rev. Lett.* **95**, 010402 (2005).
 - [54] R. Gati, M. Albiez, J. Fölling, B. Hemmerling and M.K. Oberthaler, *Appl. Phys. B* **82**, 207 (2006).
 - [55] R. Gati and M. K. Oberthaler, *J. Phys. B: At. Mol. Opt. Phys.* **40**, R61 (2007).
 - [56] B. Chatterjee, I. Brouzos, L. Cao and P. Schmelcher, *Phys. Rev. A* **85**, 013611 (2012).
 - [57] *Applied Combinatorial Mathematics*, ed. by E.F. Beckenbach, JOHN WILEY AND SONS, pp. 27–30 (1964).